arXiv:1211.5364v3 [math.AC] 1 Dec 2012

The $\mathrm{N}_{2,p}$-property of binomial extensions of simplicial complexes

Hernan de Alba Casillas, Marcel Morales

Abstract

M. Morales introduced a family of binomial ideals that are binomial extensions of square free monomial ideals. Let $I \subset \mathbb{k}[\mathbf{x}]$ be a square free monomial ideal and $J \subset \mathbb{k}[\mathbf{z}]$ a sum of scroll ideals with some extra conditions, we define the binomial extension of I as $\mathcal{B} = I + J \subset \mathbb{k}[\mathbf{z}]$. We set $p_2(\mathcal{B})$ the minimal $i \in \mathbb{N}$ such that there exists $j > 2$ such that $\beta_{i,i+j}(\mathcal{B}) \neq 0$. In the case where $J = 0$, Fröberg characterized combinatorally the case $p_2(I) = \infty$; later Eisenbud et al. solved the case $p_2(I) < \infty$. We obtain a similar result as Fröberg for the binomial extensions and we find lower and upper bounds of $p_2(\mathcal{B})$ for some families of binomial extensions in combinatorial terms as Eisenbud et al. With some additional hypothesis we can compute $p_2(\mathcal{B})$.

1 Preliminaries

$\mathbf{N}_{d,p}$-property

Let V be a set of variables and $\mathbb{k}[\mathbf{x}] = \mathbb{k}[V]$ the polynomial ring on the set of variables V over a field $\mathbb{k}$, multigraded by a commutative cancellative semigroup Σ with unity and without inversible elements excepts the unity. We simply say $\mathbb{k}[\mathbf{x}]$ is Σ-(positively) multigraded. Let $I \subset \mathbb{k}[\mathbf{x}]$ a Σ-multigraded ideal of $\mathbb{k}[\mathbf{x}]$. A Σ-multigraded minimal free resolution of I is given by:

$$0 \to \bigoplus_{\substack{j=1\\ a_{j,\rho}\in\Sigma}}^{n_\rho} \mathbb{k}[\mathbf{x}](-a_{j,\rho})^{\beta_{\rho,a_{j,\rho}}} \to \cdots \to \bigoplus_{\substack{j=1\\ a_{j,0}\in\Sigma}}^{n_0} \mathbb{k}[\mathbf{x}](-a_{j,0})^{\beta_{0,a_{j,0}}} \to I \to 0,$$

where $\beta_{i,a_j} \in \mathbb{N}^*$ and $a_j \in \Sigma$. The ranks $\beta_{i,a_j}(I) = \beta_{i,a_j}$ are called Σ-multigraded (or simply Σ-graded) Betti numbers. In case where $\Sigma = \mathbb{N}$ and $\mathbb{k}[\mathbf{x}]$ have the $\mathbb{N}$-standard graduation the $\mathbb{N}$-graded Betti numbers of I will just be called the Betti-numbers of I.

1.1 Theorem. (Upper-semicontinuity) *Let Σ be a semigroup and $I \subset \mathbb{k}[\mathbf{x}]$ be a Σ multigraded ideal, then for all monomial order $\succ$ on $\mathbb{k}[\mathbf{x}]$ the Betti numbers of I satisfy the following inequality:*

$$\beta_{i,a}(I) \leq \beta_{i,a}(\mathrm{in}_\succ(I)), \text{ for all } i \in \mathbb{N} \text{ and } a \in \Sigma.$$

A good reference for theorem 1.1 is chapter 8 of [19].

1.2 Remark. If there exists a monomial order $\succ$ such that $\beta_{i,a}(\mathrm{in}_\succ(I)) = 0$, then $\beta_{i,a}(I) = 0$.

1.3 Definition. Let $I \subset \mathbb{k}[\mathbf{x}]$ be a homogeneous ($\mathbb{N}$-graded standard) ideal of $\mathbb{k}[\mathbf{x}]$ and $d, p \in \mathbb{N}$, $d \neq 0$ and $p \neq 0$. We say that I satisfies the $\mathrm{N}_{d,p}$ property if and only if for all $i \leq p-1$ and $j \geq 1$ $\beta_{i,i+d+j}(I) = 0$. Now, if $d \in \mathbb{N}$, $d \neq 0$ we set $p_d(I) \in \mathbb{N}$ as the maximal integer p such that I satisfies the $\mathbf{N}_{d,p}$-property, if this maximum exists, otherwise we set $p_d(I) = \infty$ and in this case we say that I is d-regular. In case where I is generated by homogeneous elements of degree d and is d-regular, we say that I is d-linear.

1.4 Proposition. *Let $I \subset \Bbbk[\mathbf{x}]$ be an ideal generated by homogeneous elements of degree d, $\succ$ a monomial order over $\Bbbk[\mathbf{x}]$, then*

$$\beta_{i,j}(I) = \beta_{i,j}(\mathrm{in}(I)) \textit{ for all } i \leq p_d(\mathrm{in}(I)) - 1 \textit{ and } j \in \mathbb{N}\textit{; and}$$
$$\beta_{p_d(\mathrm{in}(I)),p_d(\mathrm{in}(I))+d}(I) = \beta_{p_d(\mathrm{in}(I)),p_d(\mathrm{in}(I))+d}(\mathrm{in}(I)),$$

thus $p_d(I) \geq p_d(\mathrm{in}(I))$. In particular, if $\mathrm{in}(I)$ is d-linear, then I is also d-linear and $\beta_{i,j}(I) = \beta_{i,j}(\mathrm{in}(I))$ for all $i, j \in \mathbb{N}$.

Proof. We set $p = p_d(\mathrm{in}(I))$. So the Betti-table of $\mathrm{in}(I)$ has the shape:

	0	1	$\dots$	$p-1$	p	$\dots$	ρ
d	$\beta_{0,d}(\mathrm{in}(I))$	$\beta_{1,d+1}(\mathrm{in}(I))$	$\dots$	$\beta_{p-1,p+d-1}(\mathrm{in}(I))$	$\beta_{p,p+d}(\mathrm{in}(I))$	$\dots$	$\beta_{\rho,\rho+d}(\mathrm{in}(I))$
$d+1$	0	0	$\dots$	0	$\beta_{p,p+d+1}(\mathrm{in}(I))$	$\dots$	$\beta_{\rho,\rho+d+1}(\mathrm{in}(I))$
$\vdots$							
m	0	0	$\dots$	0	$\beta_{p,p+m}(\mathrm{in}(I))$	$\dots$	$\beta_{\rho,\rho+m}(\mathrm{in}(I))$

By the remark 1.2 for all $i < p-1$ and $j \in \mathbb{N}$, $\beta_{i,d+i+j}(I) = 0$. Let $H(\Bbbk[\mathbf{x}]/I; t)$ be the Hilbert series of $\Bbbk[\mathbf{x}]/I$, we know that

$$\frac{\mathcal{K}(I;t)}{(1-t)^n} = H(\Bbbk[\mathbf{x}]/I;t) = H(\Bbbk[\mathbf{x}]/\mathrm{in}(I);t) = \frac{\mathcal{K}(\mathrm{in}(I);t)}{(1-t)^n}$$

where $\mathcal{K}(M;t) = \sum_{i\geq 1,\ j\geq 0}(-1)^i\beta_{i-1,j}(M)t^j$ with $M = I$, or $M = \mathrm{in}(I)$. So $\mathcal{K}(I;t) = \mathcal{K}(\mathrm{in}(I);t)$ and consequently $[\mathcal{K}(I;t)]_{p+d} = [\mathcal{K}(\mathrm{in}(I);t)]_{p+d}$. Hence

$$\sum_{i=0}^{p-1}(-1)^i\beta_{i,i+d}(I)t^{i+d} + \beta_{p,d+p}(I)t^{d+p} = \sum_{i=0}^{p-1}(-1)^i\beta_{i,i+d}(\mathrm{in}(I))t^{i+d} + \beta_{p,d+p}(\mathrm{in}(I))t^{d+p}.$$

Then, for $0 \leq i \leq p$, $\beta_{i,i+d}(I) = \beta_{i,i+d}(\mathrm{in}(I))$, and $p_d(I) \geq p_d(\mathrm{in}(I))$. ∎

1.5 Remark. *Let $I \subset \Bbbk[\mathbf{x}]$ be an ideal generated by homogeneous element of degree $\geq d$, so $\beta_{0,d} = \beta_{0,d}(\mathrm{in}(I))$. Moreover, if $\mathrm{in}(I)$ is generated by monomials of degree d then I is generated by elements of degree d.*

Clique complexes

A simplicial complex Γ over a vertex set $V(\Gamma) = \{x_1, \dots, x_n\}$ is a collection of subsets of $V(\Gamma)$ such that: for all i, the set $\{x_i\}$ is in Γ; and if $F \in \Gamma$ and $G \subset F$ then $G \in \Gamma$. An element of a simplicial complex Γ is called a **face** of Γ. The maximal faces of Γ under inclusion are called facets of Γ. The 1-skeleton Γ_1 of Γ is the subcomplex of Γ consisting of all the faces of Γ whose cardinality is ≤ 2. Let $W \subset V(\Gamma)$, the restriction of Γ over W denoted by Γ_W is the simplicial complex composed by all the faces F of Γ such that $F \subset W$. We set $x^F = \prod_{x_i \in F} x_i$. The Stanley-Reisner ideal of Γ, denoted by I_Γ, is the ideal of $\Bbbk[V(\Gamma)]$ defined by $I_\Gamma = (\mathbf{x}^F : F \notin \Gamma)$.

1.6 Theorem. (**Hochster's Formula**) *All nonzero Betti numbers $\mathbb{N}^n$-graded of I_Γ and S/I_Γ lie in squarefree degrees $\sigma = (\sigma_1, \dots, \sigma_n) \in \mathbb{N}^n$, and*

$$\beta_{i,\sigma}(I_\Gamma) = \beta_{i+1,\sigma}(S/I_\Gamma) = \dim_{\Bbbk}\widetilde{H}_{|\sigma|-i-2}(\Gamma_W;\Bbbk),$$

where $W = \{i \in \{1, \dots, n\} : \sigma_i \neq 0\}$.

We recall that a graph G is a pair of sets $(V(G), E(G))$, where $E(G)$ is a family of subsets of V of cardinality 2 called edge-set of G. The edge ideal of G is $I(G) = (x_ix_k : \{x_i, x_k\} \in E(G))$. A **clique** of G is a subset T of vertices of G such that for all $v, w \in T$ $\{v, w\} \in E(G)$, i.e. the restriction of G on the vertex subset T is a complete graph. The **clique complex** of G is the simplicial complex $\Gamma(G)$ whose faces are the cliques of G. We remark that the 1-skeleton of $\Gamma(G)$ is G. So, if Γ is a simplicial complex, then Γ is a clique complex if $\Gamma = \Gamma(\Gamma_1)$, where Γ_1 is the 1-skeleton of Γ. We have that $I_\Gamma = I(G^c)$ where G^c is the graph with vertex-set $V(G)$ and edge-set $E(G^c) = \{\{v, w\} \subset V(G) : \{v, w\} \notin E(G)\}$, so I_Γ is a square-free quadratic monomial ideal and reciprocally for any square free quadratic monomial ideal I there exists a graph G such that $I = I_{\Gamma(G)}$.

1.7 Definition. A cycle C of a graph G is a subgraph of G with vertex-set $V(C) = \{v_1, \dots, v_q\}$ and its edge-set is $E(C) = \{\{v_1, v_2\}, \{v_2, v_3\}, \dots, \{v_{q-1}, v_q\}, \{v_q, v_1\}\} \subset E(G)$. We call lenght of C to be the cardinality of $E(C)$ and we denote it as $|C|$. We will say that $C \subset \Gamma$ is a cycle of Γ, where Γ is a simplicial complex, if and only if C is a cycle of Γ_1.

We say that the cycle C of lenght > 3 of G has a **chord** if there is an edge $\{v_i, v_j\} \in (E(G) \setminus E(C))$. We say that the cycle C is minimal if $|C| > 3$ and it does not have any chord, equivalently $C = G_{V(C)}$. The graph G is called **chordal graph** if all cycle of lenght > 3 has a chord.

1.8 Proposition. *Let* $\Gamma = \Gamma(G)$ *be the clique complex of the graph* G, $I = I_\Gamma$ *and* $p \geq 1$ *a natural integer. Then:*

1. $\mathrm{p}_2(I) + 3$ *is the minimal lenght of a minimal cycle.*
2. *Besides, if* $\mathrm{p}_2 \neq \infty$, $\beta_{\mathrm{p}_2(I), \mathrm{p}_2(I)+3}(I)$ *is equal to the number of minimal cycles of lenght* $\mathrm{p}_2(I) + 3$ *of* G.

Proof.

1. The first statement is the theorem 2.1 of [8].
2. We will write $p = p_2(I)$. We define $\mathfrak{C} = \{C \subset \Gamma : C$ is a minimal cycle of lenght $p + 3\}$. By (1) $\mathfrak{C}$ is not empty, because of $p = p_2(I)$. Let $C \in \mathcal{C}$, due to the formula of Hochster:
$$\beta_{i,V(C)}(I) = \dim \widetilde{H}_{|C|-i-2}(\Gamma_{V(C)}, \Bbbk) = \dim_\Bbbk \widetilde{H}_{p+3-i-2}(C, \Bbbk) = \dim_\Bbbk \widetilde{H}_{p-i+1}(C, \Bbbk),$$
and
$$\widetilde{H}_{p-i+1}(C, \Bbbk) = \begin{cases} 1 & \text{if } p - i + 1 = 1; \\ 0 & \text{else.} \end{cases}$$
Hence $\beta_{i,\mathrm{supp}(V(C))}(I) = 0$ for $i \neq p$, and $\beta_{p,V(C)}(I) = \dim_\Bbbk \widetilde{H}_1(C, \Bbbk) = 1$. So
$$\beta_{p,p+3}(I) = \sum_{\substack{\sigma \subset V(\Gamma) \\ |\sigma| = p+3}} \beta_{p,\sigma} \geq \sum_{C \in \mathcal{C}} \beta_{p,V(C)} = |\mathcal{C}|.$$
Now we have to see $\beta_{p,p+3}(I) = |\mathcal{C}|$. Let $\sigma \subset \{1, \dots, n\}$, $|\sigma| = p+3$. Due to the formula of Hochster $\beta_{p,\sigma}(I) = \widetilde{H}_1(\Gamma_\sigma, \Bbbk)$. We suppose $\widetilde{H}_1(\Gamma_\sigma, \Bbbk) \neq 0$, we will show that Γ_σ is a minimal cycle.
   - Γ_σ is a clique complex, because Γ_σ is a sub-complex of Γ.
   - Γ_σ does not have minimal cycles of lenght $\leq p + 2$, by part 1. On the other hand by [20], pp:139-141, $\widetilde{H}_1(\Gamma_\sigma, \Bbbk) \neq 0$ is generated by minimal cycles of Γ_σ, so we conclude that Γ_σ must be a minimal cycle.

Thus $\beta_{p,p+3}(I) \leq |\mathfrak{C}|$, but before we had seen $\beta_{p,p+3}(I) \geq |\mathfrak{C}|$, hence $\beta_{p,p+3}(I) = |\mathfrak{C}|$. ■

The second statement of the proposition 1.8 was already proved differently by O. Fernandez and P. Gimenez in [9].
Also, we can see that this theorem implies the Fröberg's theorem:

1.9 Theorem. ([10]) *Let $\Gamma = \Gamma(G)$ be the clique complex of the graph G. $I = I_\Gamma$ is 2-linear if and only if G is a chordal graph.*

Linearly joined sequences and 2-regularity

Let $V =< x_1, \ldots, x_n >$ be a $\Bbbk$-vector space generated by all the variables of $\Bbbk[\mathbf{x}]$. For all $Q \subset V$, let us denote $< Q > \subset V$ as the $\Bbbk$-vector space generated by Q and $(Q) \subset \Bbbk[\mathbf{x}]$ the generated ideal by Q.

Let $l \in \mathbb{N}$ and $\mathcal{J}_1, \ldots, \mathcal{J}_l$ be an ordered sequence of ideals in $\Bbbk[\mathbf{x}]$, such that $\mathcal{J}_i = (\mathcal{M}_i, (\mathcal{Q}_i))$, for all $1 \leq i \leq l$, where $\mathcal{Q}_i \subset V$ is a $\Bbbk$-vector subspace and $\mathcal{M}_i$ is an ideal containing no linear forms. Assume that the intersection $\mathcal{J}_1 \cap \cdots \cap \mathcal{J}_l$ is not redundant. The sequence of ideals $\mathcal{J}_1, \ldots, \mathcal{J}_l$ is said to be linearly joined if for all $k = 2, \ldots, l$:

$$\mathcal{J}_k + \bigcap_{i=1}^{k-1} \mathcal{J}_l = (\mathcal{Q}_k) + \left(\bigcap_{i=1}^{k-1} \mathcal{Q}_i \right). \tag{1}$$

1.10 Remark. Let $\mathcal{J}_1, \ldots, \mathcal{J}_l$ be a linearly joined sequence of ideals in $\Bbbk[\mathbf{x}]$, then the sequence $(\mathcal{Q}_1), \ldots, (\mathcal{Q}_l)$ is also linealy joined, since:

$$(\mathcal{Q}_k) + \left(\bigcap_{i=1}^{k-1} \mathcal{Q}_i \right) \subset (\mathcal{Q}_k) + \bigcap_{i=1}^{k-1} (\mathcal{Q}_i) \subset \mathcal{J}_k + \bigcap_{i=1}^{k-1} \mathcal{J}_l = (\mathcal{Q}_k) + \left(\bigcap_{i=1}^{k-1} \mathcal{Q}_i \right).$$

1.11 Theorem. (See [8] and Theorem 3 [17]) *Let $\mathcal{J}_1, \ldots, \mathcal{J}_l$ be a sequence of prime homogeneous ideals. The following statements are equivalent:*

1. *The ideal $\mathcal{J} = \mathcal{J}_1 \cap \cdots \cap \mathcal{J}_l$ is 2-regular*
2. *For all $j \in \{1, \ldots, l\}$, $\operatorname{reg}(\mathcal{J}_i) \leq 2$ and there exist an arrangement of the sequence $\mathcal{J}_1, \ldots, \mathcal{J}_l$ such that it is linearly joined.*

The next proposition follows from corollary 3 and the proposition 1 in [17].

1.12 Proposition. *Let $(\mathcal{Q}_1), \ldots (\mathcal{Q}_l)$ be a sequence of linear ideals, such that $\mathcal{Q} := (\mathcal{Q}_1) \cap \cdots \cap (\mathcal{Q}_l)$ is not redundant. The following statements are equivalents:*

1. *The ideal $\mathcal{Q}$ is 2-linear.*
2. *There exists a permutation of the prime components of $\mathcal{Q}$, namely $(\mathcal{Q}_1), \ldots, (\mathcal{Q}_l)$, such that this sequence is linearly joined.*
3. *There exist an permutation of the prime component of $\mathcal{Q}$, we say $(\mathcal{Q}_1), \ldots, (\mathcal{Q}_l)$, and for all $i \in \{2, \ldots, l\}$, there exists $\Bbbk$-linear subspaces $< \Delta_i >, \mathcal{P}_i$, such that:*
   - *(a) $\mathcal{D}_i := \oplus_{j=i+1}^{l} < \Delta_j >$;*
   - *(b) $\mathcal{Q}_i = \mathcal{D}_i + \mathcal{P}_i$;*
   - *(c) For all $k \in \{2, \ldots, l\}$, and $j < k$, $< \Delta_j > \times \mathcal{P}_j \subset (\mathcal{P}_k)$.*

For a more detailed exposition of linearly joined sequences we suggest to read [17] and [8].

Binomial extensions of clique complexes

Let Γ be a clique complex with vertex set $V(\Gamma) = \{x_0, x_1, \ldots, x_n\}$ and $\mathcal{F}_\Gamma = \{F \text{ facet of } \Gamma\}$. We recall the construction done by M. Morales and studied with M.L. Ha in [18]: let $F \subset \Gamma$ be a facet of Γ such that $\{x_{i_0}^{(F)}, x_{i_1}^{(F)}\}, \{x_0^{(F)}, x_{i_2}^{(F)}\}, \ldots, \{x_0^{(F)}, x_{i_{m_F}}^{(F)}\}$ are **proper edges**, i.e. each edge belongs uniquely to a facet who is actually F. To each proper edge, for example $\{x_0^{(F)}, x_j^{(F)}\}$, we associate a set $Y_j^{(F)}$, which could be $\emptyset$. We define $Y^{(F)} = \cup_{j=1}^{m_F} Y_j^{(F)}$ and $\overline{F} = F \cup Y^{(F)}$. So the extension of Γ, denoted by $\overline{\Gamma}$, is the complex whose facets are $\overline{F}$ for all $F \in \mathcal{F}_\Gamma$. If $Y^{(F)} \neq \emptyset$ we define $\mathcal{I}_F$ as the ideal of 2×2 minors of the scroll matrix:

$$M_F = \left(\begin{array}{cccc|ccc|c|ccc} x_{i_0}^{(F)} & y_{i_1 1}^{(F)} & \cdots & y_{i_1 n_{i_1}}^{(F)} & y_{i_2 1}^{(F)} & \cdots & y_{i_2 n_{i_2}}^{(F)} & \cdots & y_{i_{m_F} 1}^{(F)} & \cdots & y_{i_{m_F} n_{i_{m_F}}}^{(F)} \\ y_{i_1 1}^{(F)} & y_{i_1 2}^{(F)} & \cdots & x_{i_1}^{(F)} & y_{i_2 2}^{(F)} & \cdots & x_{i_2}^{(F)} & \cdots & y_{i_{k_F} 2}^{(F)} & \cdots & x_{i_{k_F}}^{(F)} \end{array} \right),$$

otherwise $\mathcal{I}_F = 0$. Hence we define the binomial extension $\mathcal{B}_{\overline{\Gamma}} \subset \Bbbk[V(\overline{\Gamma})]$ of the simplicial complex Γ as:

$$\mathcal{B}_{\overline{\Gamma}} = \mathcal{I}_{\overline{\Gamma}} + \mathcal{J}_{\overline{\Gamma}}, \text{ where } \mathcal{J}_{\overline{\Gamma}} = \textstyle\sum_{F \in \mathcal{F}_\Gamma} \mathcal{I}_F.$$

1.13 Proposition. ([15, Proposition 1.4]) *For any facet F of Γ, let $\mathcal{J}_F = (\mathcal{I}_F, (V(\overline{\Gamma}) \setminus \overline{F})) \subset \Bbbk[V(\overline{\Gamma})]$. The primary decomposition of $\mathcal{B}_{\overline{\Gamma}}$ over $\Bbbk[V(\overline{\Gamma})]$ is: $\mathcal{B}_{\overline{\Gamma}} = \bigcap_{F \text{ facette de } \overline{\Gamma}} \mathcal{J}_F$.*

2 Gröbner basis of $\mathcal{B}_{\overline{\Gamma}}$

2.1 Proposition. Let $\mathbf{x} = \{x_1, \ldots, x_n\}$ and $\mathbf{y} = \{y_1, \ldots, y_n\}$ be two families of variables which are not necessarily distinct, but we can order $\mathbf{x} \cup \mathbf{y}$ such that $x_i > x_j$ where $1 \leq i < j \leq n$ and $x_i > y_i$ where $1 \leq i \leq n$. Let $\Bbbk[\mathbf{x} \cup \mathbf{y}]$ be the polynomial ring over $\Bbbk$ with $\mathbf{x} \cup \mathbf{y}$ as indeterminates, whose monomials are ordered by the lexicographic order and M be the following matrix over $\Bbbk[\mathbf{x} \cup \mathbf{y}]$

$$M = \left(\begin{array}{cccc} x_1 & x_2 & \ldots & x_n \\ y_1 & y_2 & \ldots & y_n \end{array} \right)$$

Let $J := I_2(M)$ be the ideal generated for all minors of 2×2 over M. Thus

1. $B = \{\underline{x_i y_j} - x_j y_i : 1 \leq i < j \leq n\}$ is a minimal system of generators and B is also a Gröbner basis of J, where the underlined term is the leading term.
2. $\mathrm{in}(J)$ is a 2-linear ideal, which implies J is a 2-linear ideal.

Proof.

1. Let $f \neq g \in B$, therefore we can write them as: $f = x_i y_j - x_j y_i$ and $g = x_k y_l - x_l y_k$, where $i < j$ and $k < l$. Hence $\mathrm{in}(f) = x_i y_j$ and $\mathrm{in}(g) = x_k y_l$. We define:

$$S(f, g) := \frac{\mathrm{lcm}(\mathrm{in}(f), \mathrm{in}(g))}{\mathrm{in}(f)} f - \frac{\mathrm{lcm}(\mathrm{in}(f), \mathrm{in}(g))}{\mathrm{in}(g)} g,$$

   where $\mathrm{lcm}(\mathrm{in}(f), \mathrm{in}(g))$ is the least common multiple of $\mathrm{in}(f)$ and $\mathrm{in}(g)$. Due to the Buchberger theorem (see theorem 1.7.4 of [1]) it is enough to prove that we can reduce $S(f, g)$ to 0 by B using the division algorithm, whenever the maximal common divisor $(\mathrm{in}(f), \mathrm{in}(g)) \neq 1$. Suppose $(\mathrm{in}(f), \mathrm{in}(g)) \neq 1$. We have four cases:

   (a) $x_i = x_k$

(b) $x_i = y_l$
(b') $y_j = x_k$
(c) $y_j = y_l$

Note that cases (b) and (b') are similars, so we will only show the cases (a),(b) and (c).

(a) Since $x_i = x_k$, we have that $y_i = y_k$ and

$$S(f,g) = -y_l x_j y_i + y_j x_l y_i.$$

We can suppose $l < j$, let $h = x_l y_j - x_j y_l \in B$, so $\text{in}(h) = x_l y_j$, and $S(f,g) = y_i h \in B$.

(b) Since $x_i = y_l$,

$$S(f,g) = -x_k x_j y_i + y_j x_l y_k,$$

and $x_k > x_l > y_l = x_i > x_j > y_j$, let $h_1 = x_l y_j - x_j y_l \in B$, so $\text{in}(h_1) = x_l y_j$, and

$$S(f,g) - y_k h_1 = y_k x_j y_l - x_k x_j y_i = y_k x_j x_i - x_k x_j y_i.$$

Moreover, $h_2 = x_k y_i - y_k x_i \in B$ with $\text{in}(h_2) = x_k y_i$, so $S(f,g) - y_k h_1 = x_j h_2$, and we have that

$$S(f,g) = y_k h_1 + x_j h_2 \in B$$

(c) Since $y_j = y_l$, so $x_j = x_l$ and

$$S(f,g) = -x_k x_j y_i + x_i x_j y_k.$$

We can suppose $i < k$, let $h = x_i y_k - x_k y_i \in B$, where $\text{in}(h) = x_i y_k$, and $S(f,g) = x_j h \in B$.

Then in all the cases we conclude that $S(f,g)$ is reduced to 0 by B, for $f, g \in B$ with $(\text{in}(f), \text{in}(g)) \neq 1$. Hence B is a Gröbner basis of J. Since all the elements of B are homogeneus quadratic binomials we have that if a leading term X divide a term Y of a binomial of B, then $X = Y$. We can remark that a leading term $x_i y_j$ $(1 \leq i < j \leq n)$ of a binomial of B is not a term of other binomial of B, due to the definition of M. Thus, B is a reduce Gröbner basis of J and since B is homogeneous, B is a minimal system of generators of J.

2. Due to (1) we have that $\text{in}_{>\text{lex}}(J) = (x_i y_j : 1 \leq i < j \leq n)$ is a quadratic square free monomial ideal thanks of the order restrictions to the variables $\mathbf{x} \cup \mathbf{y}$. Besides, we can order the generators of $\text{in}(B)$ in the following diagram:

$$\begin{array}{ccccccccccc}
 & & & & & x_1y_n & & & & & \\
 & & & & x_1y_{n-1} & & x_2y_n & & & & \\
 & & & \cdot & & & & \cdot & & & \\
 & & \cdot & & & & & & \cdot & & \\
 & \cdot & & & & & & & & \cdot & \\
 & & x_1y_4 & & & & & & & & \cdot \\
 & x_1y_3 & & x_2y_4 & & & & & & & \cdot \\
x_1y_2 & & x_2y_3 & & x_3y_4 & & & & & & x_{n-1}y_n
\end{array}$$

Now, we define $\Delta_i := \{y_i\}$ and $\mathcal{P}_i := < x_1, \ldots, x_{i-1} >$, for all $i = 2, \ldots, n$;

$$\mathcal{D}_i := \oplus_{j=i+1}^{n} < \Delta_j > = < y_{i+1}, \ldots, y_n > \text{ for all } i = 1, \ldots, n \text{ and}$$
$$\mathcal{Q}_i := \mathcal{D}_i \oplus \mathcal{P}_i = \{x_1, \ldots, x_{i-1}, y_{i+1}, \ldots, y_n\}, \text{ for all } i = 1, \ldots, n.$$

We can see that for all $k \in \{2, \dots, l\}$ and $j < k$, $< \Delta_j > \times \mathcal{P}_j \subset (\mathcal{P}_k)$. Then, owing to the proposition 1.12 $\mathcal{Q}_1, \dots, \mathcal{Q}_n$ is a linearly joined sequence and $(\mathcal{Q}) = \cap_{i=1}^{n}(\mathcal{Q}_i)$ is 2-linear. In fact, by the theorem 9 [17] we have that the generators of $(\mathcal{Q})$ are the elements of the triangle above, thus $\mathrm{in}(J) = (\mathcal{Q})$ so $\mathrm{in}_{>\mathrm{lex}}(J)$ is 2-linear. Moreover, by the proposition 1.8 we have that $\beta_{i,j}(J) = \beta_{i,j}(\mathrm{in}(J))$ for all $i, j \in \mathbb{N}$. Consequently, J is also 2-linear. ■

The fact that J is 2-linear follows also from theorem 4.1 (cf. [6]) and the next proposition:

2.2 Proposition. *Let* $\mathbf{x} = \{x_1, \dots, x_n\}$ *and* $\mathbf{y} = \{y_1, \dots, y_n\}$ *be two families of variables who are not necessarily different, but we can order* $\mathbf{x} \cup \mathbf{y}$ *such that* $x_i > x_j$ *where* $1 \leq i < j \leq n$ *and* $x_i > y_i$ *where* $1 \leq i \leq n$. *We set*

$$M = \begin{pmatrix} x_1 & x_2 & \dots & x_n \\ y_1 & y_2 & \dots & y_n \end{pmatrix}$$

Then, by applying permutations on the columns of M *we obtain a scroll matrix* M'.

Proof. For $k = 1, \dots, n$ we will define a scroll matrix M'_k, such that $M'_k = [M'_{k-1}, C'_k]$.

1. Let $M'_1 = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}$.
2. Suppose that we have already built M_k. Let $\begin{pmatrix} x_{j_k} \\ y_{j_k} \end{pmatrix}$ the k-th column of M'_k.
   (a) If y_{j_k} appears in the first line of M, then there exist $j_{k+1} \in \{1, \dots, n\}$ such that $x_{j_{k+1}} = y_{j_k}$ and we take as $k+1$-th column of M'_{k+1} the column $\begin{pmatrix} y_{j_k} \\ y_{j_{k+1}} \end{pmatrix}$.
   (b) If y_{j_k} does not appears in the first line of M, then we take the first column of M that is not in M'_k in order to be the $k+1$-th column of M'_{k+1}.

It is clear that M'_{k+1} is a scroll matrix. We define $M' = M'_n$ and we have the assertion of the proposition. ■

2.3 Notation. Let M be a matrix, we will denote for $[M]_{i,j}$ the element of M that is in the i-row and j-column.

2.4 Definition. For $i \in \{1, \dots, k\}$ let M_i be a $2 \times n_i$ matrix. We say that $M_1, \dots, M_k$ are admissibly ordered if and only if for all $1 \leq i \leq k$ one of the following statements is satisfied:

1. Either $[M_{F_i}]_{1,1}$ is not in the second row of M_{F_j}, for all $j > i$;
2. or there exists $j > i$ such that $[M_{F_i}]_{1,1} = [M_{F_j}]_{2,1}$ and there exists $i' < i$ such that $[M_{F_{i'}}]_{1,1} = [M_{F_j}]_{1,1}$ and for all $j' < i'$, $[M_{F_i}]_{1,1} \neq [M_{F_{j'}}]_{1,1}$.

2.5 Definition. Let Γ be a clique complex and $\overline{\Gamma}$ be a binomial extension of Γ. Let $\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\}$. We will say that $\mathcal{F}$ is endowed with an admissible order if there is an order for $\mathcal{F}$, namely, $\mathcal{F} = \{F_1, \dots, F_k\}$, such that the matrices $M_{F_1}, \dots, M_{F_k}$ are admissibly ordered.

2.6 Definition. Let M be a $2 \times n$ matrix, as in the proposition 2.2. Let $\Pi \in S_n$, we define

$$\Pi(M) = \begin{pmatrix} x_{\Pi(1)} & x_{\Pi(2)} & \dots & x_{\Pi(n)} \\ y_{\Pi(1)} & y_{\Pi(2)} & \dots & y_{\Pi(n)} \end{pmatrix}$$

We say that the permutation $\Pi \in S_n$ is an **admissible permutation** of M if $\Pi(1) = 1$ and for all $i \in \{2, \dots, n\}$ and $j \leq i$, $y_{\Pi(i)} \neq x_{\Pi(j)}$.

2.7 Definition. Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ and $\mathcal{F} = \{F_1, \dots, F_k\}$. Let $\mathcal{P}_{\overline{\Gamma}}$ the set:

$$\{(\Pi_1, \dots, \Pi_k) \in S_{|Y_{F_1}|+1} \oplus \cdots \oplus S_{|Y_{F_k}|+1} : \forall h \in \{1, \dots, k\}\ \Pi_h \in S_{|Y_{F_h}|+1} \text{ admissible permutation of } M_{F_h}\}.$$

2.8 Definition. Let $\Gamma(G)$ be a clique complex, $\overline{\Gamma}$ be a binomial extension of Γ and $\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\}$ is endowed with an admissible order, namely $\mathcal{F} = \{F_1, \dots, F_k\}$. For $\Pi \in \mathcal{P}_{\overline{\Gamma}}$, let $L'_1(\Pi_h(M_{F_h}))$ be the set of all the vertices that appear in the first row of the matrix M_{F_h} and does not in the first row of a matrix of a previous facet. We define an order $>_\Pi$ on the vertex set of $\overline{\Gamma}$ in the following way: $L'_1(\Pi_h(M_{F_h}))$ is ordered by the decreassing order of the columns and if $z \in L'_1(\Pi_h(M_{F_h}))$ and $z' \in L'_1(\Pi_{h'}(M_{F_{h'}}))$ so $z > z'$ if and only if $h < h'$. The remaining vertices are ordered by an arbitrary order. In case where Π_k is the identity in $S_{|Y_{F_h}|+1}$ for all $h \in \{1, \dots, k\}$, we will denote $>_\Pi$ by $>$.

2.9 Remark. Since $M_{F_1}, \dots, M_{F_k}$ are admissibly ordered by the definition 2.8, we have in this case that for all $\Pi \in \mathcal{P}$, the order $>_\Pi$ of the vertices of $\overline{\Gamma}$ satisfies that for all $1 \leq i \leq k$ and $1 \leq j < s \leq |Y^{(F_i)}| + 1$, $[\Pi_i(M_{F_i})]_{1,j} >_{\lambda_\Pi} [\Pi_i(M_{F_i})]_{1,s}$; and for all $1 \leq j < s \leq |Y^{(F_i)}| + 1$ and $1 \leq j \leq |Y^{(F_i)}| + 1$, $[\Pi_i(M_{F_i})]_{1,j} >_{\lambda_\Pi} [\Pi_i(M_{F_i})]_{2,j}$.

2.10 Definition. Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$. We define the simplicial complex $\overline{\Gamma}^{(\Pi)}$ as the clique complex generated by the graph $(\overline{\Gamma}^{(\Pi)})_1$ obtained from $(\overline{\Gamma})_1$ deleting all the edges which correspond to all the diagonals of $\Pi(M_F)$ from the top to the bottom from left to right, where F is a facet of Γ such that $F \neq \overline{F}$. If Π is the identity we will write $\overline{\Gamma}'$ instead of $\overline{\Gamma}^{(\Pi)}$.

2.11 Remark. Thanks to the definition 2.10, for any facet $F \in \Gamma$, the restriction $\overline{\Gamma}^{(\Pi)}_{\overline{F}}$ coincide with $< F >$ if $F = \overline{F}$, and coincide with $\Gamma(\text{in}(\mathcal{I}_F))$ if $F \neq \overline{F}$. As $\mathcal{I}_F$ for all $F \neq \overline{F}$ is a scroll ideal, $\mathcal{I}_F$ is 2-linear and by Fröberg's (theorem 1.9) we have that $\Gamma(\text{in}(\mathcal{I}_F))_1$ is a chordal graph. If $F = \overline{F}$, $\overline{\Gamma}^{(\Pi)}_{\overline{F}} =< F >$ is the simplex F so $\overline{\Gamma}^{(\Pi)}_{\overline{F}}$ is a chordal graph. Thus, for all F facet of Γ, $(\overline{\Gamma}^{(\Pi)}_{\overline{F}})_1$ is a chordal graph.

Let Γ be a clique complex; $\overline{\Gamma}$ a binomial extension of Γ. Set: $\text{NF}_{\overline{\Gamma}}$ the generator set of $\mathcal{I}_{\overline{\Gamma}}$, i.e.

$$\text{NF}_{\overline{\Gamma}} = \{\mathbf{x}^{\text{supp}(\sigma)} | \sigma \notin \overline{\Gamma} \text{ and } |\sigma| = 2\}$$

and J_F the set of 2×2 minors of the matrix M_F. Then $B_{\overline{\Gamma}} = \text{NF}_{\overline{\Gamma}} \bigcup (\cup_{k=1}^{l} \text{J}_{F_k})$ is a generator system of $\mathcal{B}_{\overline{\Gamma}}$.

2.12 Proposition. *Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ. We suppose $\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\}$ is endowed with an admissible order. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$. We order the vertices of $\overline{\Gamma}$ by $>_\Pi$ (see definition 2.8). Then*

1. *$B_{\overline{\Gamma}}$ is a Gröbner basis of $\mathcal{B}_{\overline{\Gamma}}$ to the lexicographic order on the monomials of $\Bbbk[\mathbf{z}]$.*
2. *The ideal $\text{in}_{>_{\Pi\text{lex}}}(\mathcal{B}_{\overline{\Gamma}})$ is a square free quadratic monomial ideal and the associated simplicial complex of $\text{in}_{>_{\Pi\text{lex}}}(\mathcal{B}_{\overline{\Gamma}})$ is $\overline{\Gamma}^{(\Pi)}$.*

Proof.

1. If $\mathcal{I}_F$ is not trivial, then $\mathcal{I}_F$ is generated by the minors of $\Pi(M_F)$, i.e.,

$$[\Pi(M_F)]_{1,i}[\Pi(M_F)]_{2,k} - [\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,i}$$

with $i < k$, where $[\Pi(M_F)]_{1,i}[\Pi(M_F)]_{2,k}$ is the initial term for the order $\succ_\Pi$.
By using the Buchberger's criteria we need to prove that

$$S(f,g) = \frac{l.c.m(\mathrm{in}(f),\mathrm{in}(g))}{\mathrm{in}(f)}f - \frac{l.c.m(\mathrm{in}(f),\mathrm{in}(g))}{\mathrm{in}(g)}g \xrightarrow{B_{\overline{\Gamma}}} 0$$

for all $f, g \in B_{\overline{\Gamma}}$ whenever $(\mathrm{lm}(f), \mathrm{lm}(g)) \neq 1$. We have four cases:

(a) $f, g \in \mathrm{NF}_{\overline{\Gamma}}$, so they are monomials and $S(f,g) = 0$

(b) $f, g \in \mathrm{J}_F$ for any facet F of Γ. As f, g are minors of the matrix M_F and $[\Pi(M_F)]_{1,i} >_{\lambda_\Pi} [\Pi(M_F)]_{1,j}$, where $1 \leq i < j \leq |Y^{(F)}| + 1$ and $[\Pi(M)]_{1,i} \succ [\Pi(M)]_{2,i}$ for $i \in \{1,\dots,|Y^{(F)}|+1\}$; thus, by the proposition 2.1 J_F is a Gröbner basis of $\mathcal{I}_F$, and then $S(f,g) \xrightarrow{J_F} 0$.

(c) $f \in \mathrm{NF}_{\overline{\Gamma}}$ and $g \in \mathrm{J}_F$ for some facet F of Γ. Let

$$g = [\Pi(M_F)]_{1,i}[\Pi(M_F)]_{2,k} - [\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,i},$$

where $i < k$, and $f = ab$. Hence, we can suppose that $a = [\Pi(M_F)]_{1,i}$ or $a = [\Pi(M_F)]_{2,k}$. So $b \notin F$ and $S(f,g) = b[\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,i}$. Besides $[\Pi(M_F)]_{1,k} \in Y^{(F)}$. Consequently $b[\Pi(M_F)]_{1,k} \in \mathrm{NF}_{\overline{\Gamma}}$ and $S(f,g) \to 0$.

(d) $f \in \mathrm{J}_F$ and $g \in \mathrm{J}_{F'}$ for two different facets F and F' of Γ. Since $(\mathrm{in}(f), \mathrm{in}(g)) \neq 1$, the matrices M_F and $M_{F'}$ must have a common vertex x with $x \in F \cap F'$ such that $x|\mathrm{in}(f)$ and $x|\mathrm{in}(g)$. We have four possibilities for this vertex x:

i. Either $x = [\Pi(M_F)]_{1,1} = [\Pi(M_{F'})]_{1,1}$.
Hence $f = x[\Pi(M_F)]_{2,k} - [\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,1}$ and $g = x[\Pi(M_{F'})]_{2,r} - [\Pi(M_{F'})]_{1,r}[\Pi(M_{F'})]_{2,1}$ for some $k \in \{1,\dots,|Y^{(F)}|+1\}$ and some $r \in \{1,\dots,|Y^{(F')}|+1\}$. Thus

$$S(f,g) = -[\Pi(M_{F'})]_{2,r}[\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,1} + [\Pi(M_F)]_{2,k}[\Pi(M_{F'})]_{1,r}[\Pi(M_{F'})]_{2,1}.$$

For $[\Pi(M_F)]_{2,k}$ we have to cases:

A. $[\Pi(M_F)]_{2,k} \in V(\Gamma)$. Then $[\Pi(M_F)]_{2,k} \notin \overline{F'}$. Otherwise $\{x, [\Pi(M_F)]_{2,k}\} \in \overline{F'}$ and $\{x, [\Pi(M_F)]_{2,k}\}$ would not be a proper edge of F.

B. $[\Pi(M_F)]_{2,k} \notin V(\Gamma)$. Then, by definition of $\mathcal{B}_{\overline{\Gamma}}$, $[\Pi(M_F)]_{2,k} \notin \overline{F'}$.

So, in any case $[\Pi(M_F)]_{2,k} \notin \overline{F'}$, and by the same way we can see that $[\Pi(M_{F'})]_{2,r} \notin \overline{F}$. Moreover, as $k, r \geq 2$ we have that $[\Pi(M_F)]_{1,k}, [\Pi(M_{F'})]_{1,r} \notin V(\Gamma)$. Consequently

$$[\Pi(M_{F'})]_{1,r}[\Pi(M_F)]_{1,k},\ [\Pi(M_F)]_{2,k}[\Pi(M_{F'})]_{1,t} \in \mathrm{NF}_{\bar{\Delta}},$$

hence: $S(f,g) \xrightarrow{\{[\Pi(M_{F'})]_{2,r}y^{(F)}_{i_1}, [\Pi(M_F)]_{2,k}y^{(F')}_{j_1 1} \in \mathrm{NF}_{\bar{\Delta}}\}} 0.$

ii. Either $x = [\Pi(M_F)]_{1,1} = [\Pi(M_{F'})]_{2,t}$, where $t \in \{1,\dots,|Y^{(F')}|+1\}$.
Moreover $f = x[\Pi(M_F)]_{2,k} - [\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,1}$ for some $k \in \{2,\dots,|Y^{(F)}|+1\}$ and $g = [\Pi(M_{F'})]_{1,r}x - [\Pi(M_{F'})]_{1,t}[\Pi(M_{F'})]_{2,r}$ for some $r \in \{1,\dots,|Y^{(F')}|+1\}$. Since $\mathrm{in}(g) = [\Pi(M_{F'})]_{1,r}x$, we have that $t \geq 2$ and

$$S(f,g) = -[\Pi(M_{F'})]_{1,r}[\Pi(M_F)]_{1,k}[\Pi(M_F)]^{(F)}_{2,1} + [\Pi(M_F)]_{2,k}[\Pi(M_{F'})]_{1,t}[\Pi(M_{F'})]_{2,r}.$$

As in the case i. we can prove that $[\Pi(M_F)]_{2,k} \notin \overline{F'}$ and $[\Pi(M_{F'})]_{1,r} \notin \overline{F}$.
Moreover, as $k, t \geq 2$ we have that $[\Pi(M_F)]_{1,k}, [\Pi(M_{F'})]_{1,t} \notin V(\Gamma)$. Therefore

$$S(f,g) \xrightarrow{[\Pi(M_{F'})]_{2,r}[\Pi(M_F)]_{1,k}, [\Pi(M_F)]_{2,k}y^{(F')}_{r,n_r}} 0.$$

   iii. Or $x = [\Pi(M_F)]_{1,t} = [\Pi(M_{F'})]_{2,t'}$, where $t \in \{2, \ldots, |Y^{(F)}| + 1\}$ and $t' \in \{2, \ldots, |Y^{(F')}| + 1\}$.
   Thus $f = [\Pi(M_F)]_{1,k}x - [\Pi(M_F)]_{1,t}[\Pi(M_F)]_{2,k}$ for some $k \in \{1, \ldots, |Y^{(F)}| + 1\}$ and
   $g = [\Pi(M_{F'})]_{1,r}x - [\Pi(M_{F'})]_{1,t'}[\Pi(M_{F'})]_{2,r}$ for some $r \in \{1, \ldots, |Y^{(F')}| + 1\}$.
   As in the last case we can prove that $t \geq 2$ and $t' \geq 2$. So

$$S(f,g) = -[\Pi(M_{F'})]_{1,r}[\Pi((M_F)]_{1,t}[\Pi(M_F)]_{2,k} + [\Pi(M_F)]_{1,k}[\Pi(M_{F'})]_{1,t'}[\Pi(M_{F'})]_{2,r}.$$

   And like in the other cases we can prove that $[\Pi(M_F)]_{1,k} \notin \overline{F'}$ and $[\Pi(M_{F'})]_{1,r} \notin \overline{F}$. Besides, since $t, t' \geq 2$ we get that $[\Pi(M_F)]_{1,t}, [\Pi(M_{F'})]_{1,t'} \notin V(\Gamma)$. Then

$$[\Pi(M_{F'})]_{1,r}[\Pi(M_F)]_{1,t},\ [\Pi(M_F)]_{1,k}x[\Pi(M_{F'})]_{1,t'} \in \mathrm{NF}_{\bar{\Delta}},$$

   and $S(f,g) \xrightarrow{[M_{F'}]_{1,r}[M_F]_{1,t},[M_F]_{1,k}x[M_{F'}]_{1,t'}} 0.$

Thus for all $f, g \in \mathcal{B}_{\overline{\Gamma}}$, $S(f,g) \xrightarrow{\mathcal{B}_{\bar{\Gamma}}}_+ 0$, and by the Buchberger algorithm we can conclude $B_{\overline{\Gamma}}^{(\Pi)}$ is a Gröbner basis of $\mathcal{B}_{\overline{\Gamma}}$.

2. As for all matrix M_F,

$$\mathrm{in}([\Pi(M_F)]_{1,i}[\Pi(M_F)]_{2,k} - [\Pi(M_F)]_{1,k}[\Pi(M_F)]_{2,i}) = [\Pi(M_F)]_{1,i}[\Pi(M_F)]_{2,k}$$

with $i < k$ and $[\Pi(M_F)]_{1,i} \neq [\Pi(M_F)]_{2,k}$, we have that $\mathrm{in}(B)$ is a square-free monomial set. And by 1. B is a Gröbner basis of $\mathcal{B}_{\Gamma}$, so $\mathcal{B}_{\overline{\Gamma}} = \mathrm{in}(B)$ is a square-free monomial ideal and by definition of $\Gamma(\mathrm{in}(\mathcal{B}_{\overline{\Gamma}}))$ we get that $\Gamma(\mathrm{in}(\mathcal{B}_{\overline{\Gamma}})) = \overline{\Gamma}^{(\Pi)}$. ■

3 Existence of admissible orders

It's not obvious that given a graph G, $\Gamma(G)$ its clique complex and $\overline{\Gamma}$ a simplicial extension of Γ, $\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\}$ is endowed with an admissible order; as we can see in the next example:

3.1 Example. Let Γ be the clique complex generated by the graph G of the figure 1. Then, $F_1 = \{a, e, b\}$, $F_2 = \{d, h, a\}$, $F_3 = \{c, g, d\}$, $F_4 = \{b, f, c\}$ are the facets of Γ. We associate to each facet the following matrices:

$$M_{F_1} = \left(\begin{array}{ccc|c} a & x & y & z \\ x & y & b & e \end{array}\right),\ M_{F_2} = \left(\begin{array}{ccc|c} d & r & s & q \\ r & s & a & h \end{array}\right),$$

$$M_{F_3} = \left(\begin{array}{cc|c} c & u & t \\ u & g & d \end{array}\right) \text{ and } M_{F_4} = \left(\begin{array}{cc|c} b & w & v \\ w & c & f \end{array}\right),$$

Note that for any possible order λ of the matrices $M_{F_1}, M_{F_2}, M_{F_3}, M_{F_4}$, namely $\lambda = M_1, M_2, M_3, M_4$, thus there exists $i \in \{2, 3, 4\}$ such that $[M_1]_{1,1}$ appears in the second row of M_i. Nevertheless if we replace M_{F2} by

$$M'_{F_2} = \left(\begin{array}{ccc|c} a & s & r & h \\ s & r & d & q \end{array}\right)$$

we have that $M_{F_1}, M_{F_4}, M_{F_3}, M'_{F_2}$ are admissibly ordered and we can consider that $\mathcal{B}_{\overline{\Gamma}}$ is the binomial extension of $I_{\Gamma'}$, where Γ' is the clique complexe generated by the graph obtained from G replacing q to h.



G
Figure 1:

3.2 Remark. Let C_n be the cycle of n-vertices $x_1, \ldots, x_n$ and Γ the clique complex generated by C_n. As

$$\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\} = \{\{x_1, x_2\}, \{x_2, x_3\}, \ldots, \{x_n, x_1\}\},$$

any facet of C_n is itself a proper edge. In order to obtain a binomial extension of C_n, for any facet $e_i = \{x_i, x_{i+1}\}$ of C_n, for all $i \in \{1, \ldots, n\}$, where $x_{n+1} = x_1$, we can choose as x_0^i either x_i or x_{i+1}. So we can suppose that for all $i = 1, \ldots, n-1$: $M_{e_i} = 0$ if $Y^{(e_i)} = 0$;

$$M_{e_i} = \begin{pmatrix} x_i & y_1^{(i)} & \ldots & y_{s_i}^{(i)} \\ y_1^{(i)} & y_2^{(i)} & \ldots & x_{i+1} \end{pmatrix},$$

if $Y^{(e_i)} \neq \emptyset$. If $s_n = 0$ then $M_{e_n} = 0$; and if $Y^{(e_n)} = 0$, then

$$M_{e_n} = \begin{pmatrix} x_1 & y_1^{(n)} & \ldots & y_{s_n}^{(n)} \\ y_1^{(n)} & y_2^{(n)} & \ldots & x_n \end{pmatrix}.$$

So the family $\mathcal{M} = \{M_{e_{j_i}} \neq 0 : j_i \in \{1, \ldots, n\}\}$ is admissibly ordered. Finally we conclude $\mathcal{F}$ is endowed by an admissible order.

In the next lines we will see that there exists another family of clique complexes Γ distinct to $\{C_n : n \in \mathbb{N}\}$ which satifies that $\mathcal{F}$ is endowed by an admissible order.

3.3 Remark. Clearly, if $|\mathcal{F}| = 1$, $\mathcal{F}$ is endowed with an admissible order. In addition, if $\mathcal{F}$ is endowed by an admissible order, then any subfamily $\mathcal{F}' \subset \mathcal{F}$ is endowed with an admissible order, because the admissible order of $\mathcal{F}$ restricted over the subfamily $\mathcal{F}' \subset \mathcal{F}$ is an admissible order of $\mathcal{F}'$.

3.4 Notation. *Let $F \neq \overline{F}$ facet of Γ, we set $L_2(M_{F_i})$ the set of all the elements of the second line of M_{F_i}.*

3.5 Lemma. *Let Γ be a clique complex and $\overline{\Gamma}$ a binomial extension of Γ. Let $\mathcal{F}'$ be an ordered subfamily, that is $\mathcal{F}' = \{F_1, \ldots, F_s\} \subset F$ where $s > 1$, such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{1, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$, then $\mathcal{F}'$ is not endowed with any admissible order.*

Proof. Let $\Pi \in S_s$ any permutation of $\{1, \ldots, s\}$, then $M_{F_{\Pi(1)}}, \ldots, M_{F_{\Pi(s)}}$ are not admissibly ordered, because:

- If $\Pi(1) \neq s$, then $[M_{F_{(\Pi(1))}}]_{1,1} \in L_2(M_{F_{\Pi(1)+1}})$, thus there exists $1 < j \leq s$ such that $\Pi(j) = \Pi(1) + 1$; and $M_{F_{\Pi(1)}}, \ldots, M_{F_{\Pi(s)}}$ are not admissibly ordered.

- If $\Pi(1) = s$, then there exists $1 < j \leq s$ such that $\Pi(j) = 1$ and
$$[M_{F_{(\Pi(1))}}]_{1,1} \in L_2(M_{F_1}) = L_2(M_{F_{\Pi(j)}});$$
$F_{\Pi(1)}, \ldots, F_{\Pi(s)}$ are not admissibly ordered.

In this manner we conclude that $\mathcal{F}'$ is not endowed with an admissible order. ■

3.6 Proposition. *Let* Γ *be a clique complex generated by* G *and* $\overline{\Gamma}$ *a binomial extension of* Γ. $\mathcal{F}$ *is not endowed with an admissible order if and only if there exists a subfamily* $\mathcal{F}'$ *such that* $\mathcal{F}'$ *has an order* $\mathcal{F}' = \{F_1, \ldots, F_s\}$ *where* $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ *for all* $i \in \{1, \ldots, s-1\}$ *and* $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$.

Proof.

$\Rightarrow$) Let $1 \leq s \leq k$ be the integer satisfying that for any subfamily $\mathcal{F}^* \subset \mathcal{F}$ with $|\mathcal{F}^*| < s$, $\mathcal{F}^*$ is endowed with an admissible order and there exists a subfamily $\mathcal{F}' \subset \mathcal{F}$ with $|\mathcal{F}'| = s$ is not endowed with an admissible order. Obviously $s > 1$. We are going to prove that for the family $\mathcal{F}'$ there exists an order $\mathcal{F}' = \{F_1, \ldots, F_s\}$ such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{1, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$.

1. Assume that there exists $M_F \in \mathcal{F}'$ such that for all $M_{F'} \in \mathcal{F}'$, $F' \neq F$,
$$[M_F]_{1,1} \notin L_2(M_{F'}).$$
By hypothesis, $\mathcal{F}' \setminus \{F\}$ is endowed with an admissible order, namely $\mathcal{F}' \setminus \{F\} = \{F_1, \ldots, F_{s-1}\}$. Then $[M_F]_{1,1} \notin L_2(M_{F_i})$ for all $i = 1 \ldots, s-1$, and in this way $M_F, M_{F_{i_1}}, \ldots, \ldots, M_{F_{i_{s-1}}}$ are admissibly ordered, but it is a contradiction because we had supposed that $\mathcal{F}'$ was not endowed with an admissible order. Consequently: $\forall F \in \mathcal{F}'$ there exists $F' \in \mathcal{F}'$, $F' \neq F$ such that $[M_F]_{1,1} \in L_2(M_{F'})$.
2. Now we are going to prove that for the family $\mathcal{F}'$ there exists an order $\mathcal{F}' = \{F_1, \ldots, F_s\}$ such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{1, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$.
Let any $F_1 \in \mathcal{F}'$, by the point 1, there exists $F_2 \neq F_1$ such that $[M_{F_1}]_{1,1} \in L_2(M_{F_2})$. Moreover by the same point 1, there exists $F \in (\mathcal{F}' \setminus F_2)$ such that $[M_{F_2}]_{1,1} \in L_2(M_F)$.
   - If $F = F_1$, then the family of facets $\{F_1, F_2\}$ is not endowed with an admissible order, and by the lemma 3.5 and by maximality of s, $s = 2$ and $\mathcal{F}' = \{F_1, F_2\}$ satisfies that $[M_{F_i}]_{1,1} \in L_2(M_{F_2})$ and $[M_{F_2}]_{1,1} \in L_2(M_{F_1})$.
   - If $F \neq F_1$, $s > 2$, we call $F_3 := F$. Thanks to 1. there exists $F \in \mathcal{F}' \setminus F_3$ such that $[M_{F_2}]_{1,1} \in L_2(M_{F_3})$.
     - $*$ If $F = F_1$, then the family of facets $\{F_1, F_2, F_3\}$ is not endowed with an admissible order, and by the lemma 3.5 and by maximality of s, $s = 3$ and the family $\mathcal{F}' = \{F_1, F_2, F_3\}$ is endowed by an admissible order that we wanted to prove.
     - $*$ If $F = F_2$, the subfamily of facets $\{F_2, F_3\}$ does not allow and admissible order by the lemma 3.5 but it is not possible because $s > 2$ is maximal.
     - $*$ Thus if $F \neq F_1$, then $F \neq F_2$ and $s > 3$. We set $F_4 = F$ and by induction we can suppose that we have found a sequence of differents facets in $\mathcal{F}'$: $F_1, F_2, F_3, \ldots, F_s$ such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i = 1, \ldots, s-1$. Thanks to item 1. there exists $F \in \mathcal{F}' \setminus F_{s-1}$ such that $[M_{F_1}]_{1,1} \in L_2(M_F)$. If $F \neq F_1$, we can say $F = F_j$ with $j > 2$, then $F_j, F_{j+1}, \ldots, F_s$ satisfies that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{j, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_j})$. By the lemma 3.5 the family $\mathcal{F}'' = \{F_j, F_{j+1}, \ldots, F_s\}$ is not endowed with an admissible order and it is a contradiction to the maximality of s. Then $F = F_1$, and $\mathcal{F}'$ satisfies that there exists an order $\mathcal{F}' = \{F_1, \ldots, F_s\}$ such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{1, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$.

$\Leftarrow$) By the lemma 3.5 $\mathcal{F}'$ is not endowed with an admissible order. Thus, by the remark 3.3, $\mathcal{F}$ is not endowed with an admissible order. ■

3.7 Lemma. *Let G be a graph, $\Gamma := \Gamma(G)$ the clique complex generated by G and $\overline{\Gamma}$ a binomial extension of Γ. If $|\mathcal{F}| \leq 3$, then $\mathcal{F}$ is endowed with an admissible order.*

Proof.

- $\mathbf{k=2}$. If the family $\mathcal{F}$ doesn't allow any admissible order, by the proposition 3.6 there exist an order of $\mathcal{F}$: $\mathcal{F} = \{F_1, F_2\}$ such that $[M_{F_1}]_{1,1} \in L_2(M_{F_2})$ and $[M_{F_2}]_{1,1} \in L_2(M_{F_1})$. Thus $\{[M_{F_1}]_{1,1}, [M_{F_2}]_{1,1}\} \in F_1 \cap F_2$ and $\{[M_{F_1}]_{1,1}, [M_{F_2}]_{1,1}\}$ is not a proper edge of Γ, but it is not possible because $\overline{\Gamma}$ is a binomial extension of Γ. Consequently $\mathcal{F}$ is endowed with an admissible order.
- $\mathbf{k=3}$. If the family $\mathcal{F}$ is not endowed with an admissible order, then by the proposition 3.6 there exist an order of $\mathcal{F}$: $\mathcal{F} = \{F_1, F_2, F_3\}$ such that $[M_{F_1}]_{1,1} \in L_2(M_{F_2})$, $[M_{F_2}]_{1,1} \in L_2(M_{F_3})$ and $[M_{F_3}]_{1,1} \in L_2(M_{F_1})$. So we can conclude that $e_1 = \{[M_{F_2}]_{1,1}, [M_{F_1}]_{1,1}\} \in F_2$, $e_2 = \{[M_{F_3}]_{1,1}, [M_{F_2}]_{1,1}\} \in F_3$, $e_3 = \{[M_{F_1}]_{1,1}, [M_{F_3}]_{1,1}\} \in F_1$ and $L = \{[M_{F_1}]_{1,1}, [M_{F_2}]_{1,1}, [M_{F_3}]_{1,1}\}$ is a clique of G, thus $L \subset F_i$ for some $i \in \{1,2,3\}$ and in this way either e_1, either e_2 or e_3 is not a proper edge, and it is a contradiction to the fact that $\overline{\Gamma}$ is a binomial extension of Γ. Thus, we conclude that $\mathcal{F}$ is endowed with an admissible order. ■

3.8 Proposition. *Let G be a graph, $\Gamma := \Gamma(G)$ the clique complex generated by G and $\overline{\Gamma}$ a binomial extension of Γ. Assuming that the graph G restricted on the family $\mathcal{F}$ is a chordal graph, then $\mathcal{F}$ is endowed with an admissible order.*

Proof. If $|\mathcal{F}| = 1, 2$ or 3 by the lemma 3.7 the family $\mathcal{F}$ is endowed by an admissible order.

Let us suppose that $|\mathcal{F}| \geq 4$. Assume that the family $\mathcal{F}$ is not endowed with an admissible order. So, by the proposition 3.6 and lemma 3.7 there exists a subfamily $\mathcal{F}' = \{F_1, \ldots, F_s\} \subset \mathcal{F}$ where $s > 3$, such that $[M_{F_i}]_{1,1} \in L_2(M_{F_{i+1}})$ for all $i \in \{1, \ldots, s-1\}$ and $[M_{F_s}]_{1,1} \in L_2(M_{F_1})$. Thus, by definition of binomial extension, $\{[M_{F_{i+1}}]_{1,1}, [M_{F_i}]_{1,1}\}$ is a proper edge of F_{i+1} for all $i \in \{1, \ldots, s-1\}$ and $\{[M_{F_s}]_{1,1}, [M_{F_1}]_{1,1}\}$ is a proper edge of F_1. So, $\{\{[M_{F_1}]_{1,1}, [M_{F_2}]_{1,1}\}, \ldots, \{[M_{F_s}]_{1,1}, [M_{F_1}]_{1,1}\}\}$ are the edges of a cycle C of G, but since G restricted on the family $\mathcal{F}$ is a chordal graph, C is not a minimal cycle and C must have a clique $V \subset V(C)$ of size 3 with two edges e, e' of C. In this way we have that $V \subset F_i$ for some $i \in \{1, \ldots, k\}$ and at least an edge among e and e' is in two facets of Γ, but this is a contradiction to the fact that e, e' are proper edges of Γ. Consequently $\mathcal{F}$ is endowed with an admissible order. ■

4 Lower bounds of $p_2(\mathcal{B}_{\overline{\Gamma}})$

Let Γ be a clique complex and $\overline{\Gamma}$ be a binomial extension of Γ. We suppose that $\mathcal{F} = \{F$ facet of Γ : $\overline{F} \neq F\}$ is endowed with an admissible order, namely $\mathcal{F} = \{F_1, \ldots, F_k\}$. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$ (see definition 2.7). Using the definition 2.8 we get and order $>_\Pi$ for the vertex-set of $\overline{\Gamma}$ and by the proposition 2.12 the associated simplicial complex of $\mathrm{in}_{>_\Pi \mathrm{lex}}(B_{\overline{\Gamma}})$ is $\overline{\Gamma}^{(\Pi)}$ defined in 2.10. Using the proposition 1.4 we have that

$$\max\nolimits_{\Pi \in \mathcal{P}_{\overline{\Gamma}}}(p_2(I(\overline{\Gamma}^{(\Pi)}))) \leq p_2(B_{\overline{\Gamma}}). \tag{2}$$

And from proposition 1.8, let $\mathfrak{C}'_\Pi$ be the family of all minimal cycles of $\overline{\Gamma}^\Pi$

$$\max_{\Pi\in\mathcal{P}_{\overline{\Gamma}}}(p_2(I(\overline{\Gamma}^{(\Pi)}))) + 3 = \max_{\Pi\in\mathcal{P}_{\overline{\Gamma}}}(min_{C'\in\mathfrak{C}'_\Pi}|C'|) \tag{3}$$

So we would like to know the family of $\mathfrak{C}'_\Pi$ for any $\Pi \in \mathcal{P}_{\overline{\Gamma}}$ in order to calculate $min_{C'\in\mathfrak{C}'_\Pi}|C'|$.

4.1 Definition. Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ, $\mathcal{F} = \{F_1, \ldots, F_k\}$ and $\Pi \in \mathcal{P}_{\overline{\Gamma}}$. We say that a cycle C of Γ is a **Π-virtual minimal cycle** if and only if either C is a minimal cycle of Γ, or C is not a minimal cycle in Γ, but C satisfies the following properties:

1. Any chord of C is not an edge of $(\overline{\Gamma}^{(\Pi)})_1$.
2. If $e, e' \in E(C)$ are distinct, then e, e' are not in a same facet of Γ.

4.2 Remark. Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ and $\Pi, \Pi' \in \mathcal{P}_{\overline{\Gamma}}$. If $e \in (E(G) \setminus E(\overline{G}^{(\Pi)}))$, then $e \in (E(G) \setminus E(\overline{G}^{(\Pi')}))$. So the definition of a Π-virtual minimal cycle of G does not depend on Π, so a Π-virtual minimal cycle can just be called **virtual minimal cycle**. We denote by $\mathfrak{C}^v$ the family of all the virtual minimal cycles of Γ.

4.3 Proposition. *Let Γ be a clique complex and $\overline{\Gamma}$ be a binomial extension of Γ. Then, for any $C \in \mathfrak{C}^v$ there exists $V \subset V(C)$ such that Γ_V is a minimal cycle of lenght ≥ 4.*

Proof. Asume that $E(C) = \{\{x_1, x_2\}, \{x_2, x_3\}, \ldots, \{x_s, x_1\}\}$. Let consider the sets $V \subset V(C)$ such that Γ_V contains a cycle. If for any of them all the cycles contained in Γ_V are not minimal cycles. So, there exist $j \in \{1, \ldots, s\}$ such that $\{z_j, z_{j+2}\}$ is a chord of C, i.e. $D = \{z_j, z_{j+1}, z_{j+2}\}$ is a clique of Γ, so there exists a facet F in Γ who contains D in contradiction with the item 2 of the definition 4.1. Thus, there exists $V \subset V(C)$ such that Γ_V is a minimal cycle of Γ. ■

4.4 Definition. Let Γ be the clique complex and $\overline{\Gamma}$ be a binomial extension of Γ. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$, C be a minimal virtual cycle of Γ and $e \in E(C)$ and $e = \{x_e, x'_e\}$ be an edge in $E(C)$. We say e is **virtual** if $e \notin E(\overline{\Gamma}^{(\Pi)})$. By the remark 4.2 it does not depend on Π, i.e. if $e \notin E(\overline{\Gamma}^{(\Pi)})$, then $e \notin E(\overline{\Gamma}^{(\Pi')})$, for any $\Pi' \in \mathcal{P}_{\overline{\Gamma}}$. Let e be a virtual edge of C, so there exists a unique facet F_e of Γ containing e. A path P_e from x_e to x'_e in $\overline{G}^{(\Pi)}$ whose vertices are in $\overline{F}_e$ is called a **Π-local substitution** of e in $\overline{G}^{(\Pi)}$ if there is not an edge in $\overline{G}^{(\Pi)}$ such that it is a chord of P_e and for any $x \in (V(C) \setminus e)$ there is not any edge in $E(\overline{G}^{(\Pi)})$ from x to any vertex in $V(P_e) \setminus e$.

4.5 Definition. Let Γ be a clique complex, $\overline{\Gamma}$ be a binomial extension of Γ and $\mathfrak{C}$ be the family of all virtual minimal cycles of G. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$ and C be a virtual minimal cycle. We set $\mathfrak{C}^{\mathrm{ls}}_\Pi(C)$ the family of all minimal cycles C' in $\overline{G}^{(\Pi)}$ obtained from C by replacing all the virtual edges of C by local substitution in $\overline{G}^{(\Pi)}$. We set $\mathfrak{C}^{\mathrm{ls}}_\Pi = \cup_{C\in\mathfrak{C}^v}\mathfrak{C}^{\mathrm{ls}}_\Pi(C)$.

4.6 Proposition. (**Going down for minimal cycles**) *Let Γ be a clique complex and $\overline{\Gamma}$ be a binomial extension of Γ. Then, for any minimal cycle C' of lenght ≥ 4 of $\overline{\Gamma}^{(\Pi)}$, there exists a virtual minimal cycle C of Γ and a subset $V' \subset V(\overline{\Gamma})$, such that $C' = (\overline{\Gamma}^{(\Pi)})_{V(C)\cup V'}$.*

Proof. Let C' be a minimal cycle of lenght ≥ 4 of $(\overline{\Gamma}^{(\Pi)})_1$. Set $q := |C'|$,

$$E(C') = \{\{z_1, z_2\}, \{z_2, z_3\}, \ldots, \{z_q, z_1\}\},$$

and for all $i = 1, \ldots, q-1$, $e_i = \{z_i, z_{i+1}\}$ and $e_q = \{z_q, z_1\}$. For all $i \in \{1, \ldots, q\}$, there exists a facet F of Γ such that $e_i \subset \overline{F}$. We assert that there exist at least two facets $F \neq F'$ of Γ and two edges $e_i \neq e_j \in E(C')$ such that $e_i \subset \overline{F}$ and $e_j \subset \overline{F'}$, otherwise if there exist F facet of Γ such that for all edge $e \in E(C)$, $e \in \overline{F}$, so C is a subgraph of $(\overline{\Gamma}^{(\Pi)}_{\overline{F}})_1$ and by the remark 2.11, $(\overline{\Gamma}^{(\Pi)}_{\overline{F}})_1$ is a chordal graph, thus C' has a chord in $\overline{\Gamma}^{(\Pi)}$ and it is a contradition to the fact that C' is a minimal cycle of $\overline{\Gamma}^{(\Pi)}$.

Thus, we have a sequence of facets $F_{i_1}, \ldots, F_{i_q}$ in Γ such that $e_j \subset \overline{F}_{i_j}$. If $e_{j+1} \subset \overline{F}_{i_j}$, we take $F_{i_{j+1}} = F_{i_j}$. Consequently we can rewrite this sequence like $F_{i_1}, \ldots, F_{i_{s'}}$, where $q \geq s' \geq 2$ and $F_{i_j} \neq F_{i_{j+1}}$, such that C' is divided by consecutive paths $C_j \subset \overline{F}_{i_j}$ from z_{i_j} to $z_{i_{j+1}}$, where $z_1 = z_{i_1} = z_{i_{s'+1}}$. Moreover, we can suppose that $F_{i_1} \neq F_{i_{s'}}$; since if $F_{i_1} = F_{i_{s'}}$ we can take as initial point of C' the point $z_{i_{s'}}$.

- As for all $j \in \{2, \ldots, s'+1\}$ $z_{i_j} \in \overline{F}_{j-1} \cap \overline{F}_j$, by the definition of $\overline{\Gamma}$, we have that for all $j \in \{1, \ldots, s'\}$, $z_{i_j} \in \Gamma$.
- We have that $F_{i_j} = F_{i_r}$ for any $j, r \in \mathbb{N}$ such that $r \geq j+2$. Otherwise, since C' is a minimal cycle

  $$\{z_{i_j}, z_{i_r}\}, \{z_{i_j}, z_{i_{r+1}}\}, \{z_{i_{j+1}}, z_{i_r}\}, \{z_{i_{j+1}}, z_{i_{r+1}}\} \notin E(\overline{\Gamma}^{(\Pi)})_1 \text{ (see figure 2),}$$

  but these edges are contained in F_{i_j} .



Figure 2:

  As a result of the definition of $\overline{\Gamma}^{(\Pi)}$, if $e \in (E(\Gamma) \setminus E(\overline{\Gamma}^{(\Pi)}))$ then a vertex of e is in the first line of a matrix $M_{F_{i_j}}$; thus, there exist two elements of $\{z_{i_j}, z_{i_{j+1}}, z_{i_r}, z_{i_{r+1}}\} \subset V(\Gamma)$ which appear in the first line of $M_{F_{i_j}}$. It is a contradiction with the definition of $M_{F_{i_j}}$.

Then, we have seen that there exist $s' \geq 3$ and a sequence $F_{i_1}, \ldots, F_{i_{s'}}$ such that for all $j \neq j' \in \{1, \ldots, s\}$, $F_{i_j} \neq F_{i_{j'}}$ and C' is divided in consecutive paths $C_j \subset \overline{F}_{i_j}$ from z_{i_j} to $z_{i_{j+1}}$, where $z_1 = z_{i_1} = z_{i_{s'+1}}$. Set s the smallest integer who satisfies these properties.
Let C be the cycle defined by

$$V(C) = \{z_{i_1}, z_{i_2}, \ldots, z_{i_s}\} \text{ and } E(C) = \{\{z_{i_1}, z_{i_2}\}, \ldots, \{z_{i_{s-1}}, z_{i_s}\}, \{z_{i_s}, z_1\}\}.$$

If C is a minimal cycle, then C' comes from C which is a Π-virtual minimal cycle by definition 4.1. If C is not a minimal cycle, we remark that C is a Π-virtual minimal cycle:

1. Any chord of C is not an edge in $E(\overline{\Gamma}^{(\Pi)})$, since C' is a minimal cycle of $(\overline{\Gamma}^{(\Pi)})$.
2. By construction of C and by the minimality of s, for all pair $e, e' \in E(C)$, e and e' does not belong to the same facet of Γ.

Finally, by the last three points C is a Π-virtual minimal cycle. ■

4.7 Theorem. *Let* Γ *be a clique complex and* $\overline{\Gamma}$ *a binomial extension of* Γ. *Then* Γ_1 *is a chordal graph if and only if* $\mathcal{B}_{\overline{\Gamma}}$ *is* 2*-linear.*

Proof.

$\Rightarrow$) As Γ_1 is a chordal graph, Γ does not have minimal cycles. So $(\overline{\Gamma}^{(\Pi)})$ does not too, otherwise, by the proposition 4.6 Γ would have a Π-virtual minimal cycle and by the proposition 4.3 there exists $V \subset V(C)$ such that Γ_V is a minimal cycle of Γ, but it is not possible, thus $(\overline{\Gamma}^{(\Pi)})$ does not have minimal cycles and by the proposition 1.8 we obtain that $I_{\overline{\Gamma}^{(\Pi)}}$ is 2-linear. Then, applying the proposition 1.4, we conclude that $\mathcal{B}_{\overline{\Gamma}}$ is 2-linear.

$\Leftarrow$) Assume that Γ_1 is not chordal. So there exist a minimal cycle C of Γ of lenght > 3. As $\overline{\Gamma}$ is an extension of Γ, all minimal cycle of G is a minimal cycle of $\overline{\Gamma}$, in particular C is a minimal cycle of $\overline{\Gamma}$, then by the proposition 1.8, $\mathcal{I}_{\overline{\Gamma}}$ is not 2-linear. Moreover by the proposition 1.13

$$\mathcal{B}_{\overline{\Gamma}} = \bigcap_{F \in \mathcal{F}} \mathcal{J}_F,$$

where $\mathcal{J}_F = (\mathcal{I}_F, (V(\overline{\Gamma}) \setminus \overline{F}))$. But the primary decomposition of $\mathcal{I}_{\overline{\Gamma}}$ is:

$$\mathcal{I}_{\overline{\Gamma}} = \bigcap_{F \in \mathcal{F}} (V(\overline{\Gamma}) \setminus \overline{F}),$$

thus by proposition 1.12 there does not exist a permutation to the sequence of the ideals $(V(\overline{\Gamma}) \setminus \overline{F})$ such that it is linearly joined, in this way by the remark 1.10 there does not exist a permutation of the sequence

$$\{(\mathcal{I}_F, (V(\overline{\Gamma}) \setminus \overline{F}))\}_{F \in \mathcal{F}}$$

such that it is linearly joined and so $\mathcal{B}_{\overline{\Gamma}}$ is not 2-linear by theorem 1.11. ■

4.8 Remark. Thanks to proposition 4.6 for any minimal cycle C' of $\overline{G}^{(\Pi)}$ there exist a virtual minimal cycle C of G, such that $C' \in \mathfrak{C}^{\mathrm{ls}}_{\Pi}(C)$. So, setting $\mathfrak{C}'_{\Pi}$ the family of all minimal cycles of $\overline{\Gamma}^{(\Pi)}$, then

$$\mathfrak{C}'_{\Pi} \subset \mathfrak{C}^{\mathrm{ls}}_{\Pi} \tag{4}$$

Nevertheless, it is not true in general that $\mathfrak{C}^{\mathrm{ls}}_{\Pi} = \mathfrak{C}'_{\Pi}$, as we can see in the next example:

4.9 Example. Let G be the graph of the figure 3 and $\Gamma := \Gamma(G)$ the clique complex generated by G. For the facets $F_1 = \{a, d, h\}$, $F_2 = \{a, b, e\}$, $F_3 = \{b, c, f\}$, $F_4 = \{c, d, g\}$ we associate the matrices:

$$M_{F_1} = \begin{pmatrix} a & y_1 & y_2 \\ y_1 & y_2 & d \end{pmatrix}, \; M_{F_2} = \begin{pmatrix} a & y_3 & y_4 \\ y_3 & y_4 & b \end{pmatrix},$$

$$M_{F_3} = \begin{pmatrix} b & y_5 & y_6 \\ y_5 & y_6 & c \end{pmatrix} \text{ and } M_{F_4} = \begin{pmatrix} c & y_7 & y_8 \\ y_7 & y_8 & d \end{pmatrix},$$

respectively. $\overline{G}' = (\overline{\Gamma}')_1$ as in the figure 3. We have that the cycle C defined by $E(C) = \{\{a, b\}, \{b, c\}, \{c, d\}, \{d, a\}\}$ is a minimal cycle of G, then C is a virtual cycle of G. The paths: P_1, P_2, P_3, P_4 defined by: $E(P_1) = \{\{a, e\}, \{e, b\}\}$, $E(P_2) = \{\{b, f\}, \{f, c\}\}$, $E(P_3) = \{\{c, g\}, \{g, d\}\}$, $E(P_4) = \{\{d, h\}, \{h, a\}\}$ are local substitutions of $\{a, b\}$, $\{b, c\}$, $\{c, d\}$, $\{d, a\}$, respectively, in $\overline{G}'$ but C' defined by the union of these paths is a cycle of $\overline{G}'$ but it is not a minimal cycle of $\overline{G}'$.



Figure 3:

□

Now, for any $\Pi \in \mathcal{P}_{\overline{\Gamma}}$, we would like to calculate $\min_{C' \in \mathfrak{C}^{\mathrm{ls}}_{\Pi}}(|C'|)$. The next proposition follows immediately from the inequalities 2, 3 and the remark 4.8.

4.10 Proposition. *Let* Γ *be a clique complex,* $\overline{\Gamma}$ *a binomial extension of* Γ *and* $\mathcal{F} = \{F \text{ facet of } \Gamma : F \neq \overline{F}\}$ *is endowed with an admissible order. Then*

$$p_2(\mathcal{B}_{\overline{\Gamma}}) \geq \max_{\Pi \in \mathcal{P}}(\min_{C' \in \mathfrak{C}'_{\Pi}}(|C'|)) \geq \max_{\Pi \in \mathcal{P}}(\min_{C' \in \mathfrak{C}^{\mathrm{ls}}_{\Pi}}(|C'|)).$$

■

4.11 Definition. Let G be a graph, $\Gamma = \Gamma(G)$ the clique complex generated by G and $\overline{\Gamma}$ a binomial extension of Γ. Let $\Pi \in \mathcal{P}_{\overline{\Gamma}}$ and e a virtual edge of G. A local substitution of e is called a replacement of e in $\overline{G}^{(\Pi)}$ if the lenght of the local substitution is minimal. The lenght of a replacement of e is unique and it is denoted by $t_{\Pi}(e)$. If e is not virtual we set $t_{\Pi}(e) := 1$ and in this case by notation abuse we will say that the path $P_e = \{e\}$ is a replacement of e.

4.12 Notation. *Let* F *be a facet of* Γ *such that* $F \neq \overline{F}$*. We denote by* $V_{\Gamma}(M_F)$ *the set of all the vertices in* F *that are in* M_F*.*

4.13 Proposition. *Let* Γ *be a clique complex and* $\overline{\Gamma}$ *a binomial extension of* Γ*. Let* $\Pi \in \mathcal{P}_{\Gamma}$*. Let* $C \in \mathfrak{C}^{v}$ *and* $e \in E(\Gamma)$ *be a virtual edge in* $E(C)$*, such that* $e = \{x^{(F)}_0, x^{(F)}_k\}$ *for some facet of* F *with* $Y^F \neq \emptyset$*. Then:*

1. *If there exist* $x \in (F \setminus V_{\Gamma}(M_F))$ *such that for any vertex* $x' \in V(C)$, $\{x, x'\} \notin E(\Gamma)$*, then* $t_{\Pi}(e) = 2$*.*
2. *Else, let* $J^{\mathrm{ls}}_F = \{k\} \cup \{j \in \{1, \dots, m_F\} : \forall x' \in (V(C) \setminus e),\ \{x^{(F)}_j, x'\} \notin E(\Gamma)\}$*:*
    - *(a) If* $\min_{j \in J^{\mathrm{ls}}_F}(|Y^{(F)}_j|) < |Y^{(F)}_k|$*, then* $t_{\Pi}(e) \leq \min_{1 \leq j \leq m_F}(|Y^{(F)}_j|) + 2$*.*
    - *(b) If* $\min_{j \in J^{\mathrm{ls}}_F}(|Y^{(F)}_j|) = |Y^{(F)}_k|$*, then* $t_{\Pi}(e) \leq \min_{1 \leq j \leq m_F}(|Y^{(F)}_j|) + 1$*.*

Proof.

1. If there exist $x \in F \setminus V_{\Gamma}(M_F) \neq \emptyset$ such that for any vertex $x' \in V(C)$, $\{x, x'\} \notin E(\Gamma)$, then $\{x^F_0, x\}, \{x, x^F_k\} \in E(\overline{G}^{(\Pi)})$ and $\{x^F_0, x\}, \{x, x^F_k\}$ is a replacement of e in $\overline{G}^{(\Pi)}$ and $t_{\Pi}(e) = 2$.

2. We can suppose that M_F has one of the following representations: either

$$M_F = \left(\begin{array}{cccc|ccc|c|ccc} x_0^{(F)} & y_{11}^{(F)} & \cdots & y_{1n_1}^{(F)} & y_{21}^{(F)} & \cdots & y_{2n_2}^{(F)} & \cdots & y_{m_F1}^{(F)} & \cdots & y_{m_Fn_{m_F}}^{(F)} \\ y_{11}^{(F)} & y_{12}^{(F)} & \cdots & x_1^{(F)} & y_{22}^{(F)} & \cdots & x_2^{(F)} & \cdots & y_{m_F2}^{(F)} & \cdots & x_{m_F}^{(F)} \end{array} \right), \text{ or}$$

$$M_F = \left(\begin{array}{c|ccc|c|ccc} x_0^{(F)} & y_{21}^{(F)} & \cdots & y_{2n_2}^{(F)} & \cdots & y_{m_F1}^{(F)} & \cdots & y_{m_Fn_{m_F}}^{(F)} \\ x_1^{(F)} & y_{22}^{(F)} & \cdots & x_2^{(F)} & \cdots & y_{m_F2}^{(F)} & \cdots & x_{m_F}^{(F)} \end{array} \right),$$

The following argument is valid for both representations of the matrices M_F. From now to the end of the proof we write z instead of $z^{(F)}$.

As x_0, y_{j1}, with $j \in \{2, \ldots, m_F\}$ are located only in the first line of M_F and $y_{1,1}$ appears only in the same line and the same column that x_0 in M_F, we have that for all $j \in \{1, \ldots, m_F\}$, $\{x_0, y_{j1}\}$ is always an edge of $\overline{G}^{(\Pi)}$. As $\Pi \in \mathcal{P}$ (see definition 2.6) for all $j \in \{1, \ldots, m_F\}$ and $i \in \{1, \ldots, n_j\}$, $y_{ji}y_{ji+1}$ neither $y_{jn_j}x_j$ are not diagonals of $\Pi(M_F)$ from the top to the bottom from left to right, so $\{y_{ji}, y_{ji+1}\}, \{y_{jn_j}, x_j\} \in E(\overline{G}^{(\Pi)})$. Finally, since x_j for all $j \in \{1, \ldots, m_F\}$ is only in the second line of M_F, we have that for all $j \in \{1, \ldots, k-1, k+1, \ldots, m_F\}$, $\{x_k, x_j\} \in \overline{G}^{(\Pi)}$ and $\{x_0, x_1\} \in \overline{G}^{(\Pi)}$, if $Y_1^{(F)} = \emptyset$; but if $j \in (\{1, \ldots, k-1, k+1, \ldots, m_F\} \setminus J_F^{\text{ls}})$, x_j is not a vertex of a local substitution of e. Then for all $j \in J_F^{\text{ls}}$ we can build a local substitution P_j from x_e to x'_e:

- $E(P_j) = \{\{x_0, y_{i1}\}, \{y_{i1}, y_{i2}\}, \ldots, \{y_{in_i}, x_j\}, \{x_j, x_k\}\}$ for all $j \neq k$, and
- $E(P_k) = \{\{x_0, y_{k1}\}, \{y_{k1}, y_{k2}\}, \ldots, \{y_{kn_k}, x_k\}\}$, if $j = k$.

Thus:

(a) If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) < |Y_k^{(F)}|$, then there exist $j \in J_F^{\text{ls}}$, such that P_j is a local substitution of e and $|P_j| \leq |P_{j'}|$, for all $j' \in J_F^{\text{ls}}$ so $t_\Pi(e) \leq |P_j| = \min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) + 2$.

(b) If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) = |Y_k^{(F)}|$, then $|P_k| \leq |P_j|$ with $j \in J_F^{\text{ls}}$ and

$$t_\Pi(e) \leq |P_k| = |Y_k^{(F)}| + 1 = \min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) + 1.$$

■

We recall that G is a graph, $\Gamma = \Gamma(G)$ is the clique complex generated by G and $\overline{\Gamma}$ a binomial extension of Γ. For each facet F de Γ such that $F \neq \overline{F}$, there is a matrix

$$M_F = \left(\begin{array}{cccc|ccc|c|ccc} x_0^{(F)} & y_{11}^{(F)} & \cdots & y_{1n_1}^{(F)} & y_{21}^{(F)} & \cdots & y_{2n_2}^{(F)} & \cdots & y_{m_F1}^{(F)} & \cdots & y_{i_{m_F}n_{m_F}}^{(F)} \\ y_{11}^{(F)} & y_{12}^{(F)} & \cdots & x_1^{(F)} & y_{22}^{(F)} & \cdots & x_2^{(F)} & \cdots & y_{m_F2}^{(F)} & \cdots & x_{m_F}^{(F)} \end{array} \right).$$

Actually we can write M_F like:

$$M_F = (B_0^{(F)m} | B_1^{(F)} | \cdots | B_{m_F}^{(F)}),$$

where

$$B_0^{(F)} = \begin{pmatrix} x_{i_0}^{(F)} \\ y_{i_11}^{(F)} \end{pmatrix} \text{ and } B_j^{(F)} = \begin{pmatrix} y_{i_j1}^{(F)} & \cdots & y_{i_jn_{j_1}}^{(F)} \\ y_{i_j2}^{(F)} & \cdots & x_{i_j}^{(F)} \end{pmatrix},$$

for all $j = 1, \ldots, m_F$.

4.14 Definition. Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ and F a facet of Γ different to $\overline{F}$. Let $s = \max\{n_1, \dots, n_{m_F}\}$, we denote by $\Pi^*_F(M_F)$ the column permutation of the matrix M_F such that:

$$\Pi^*_F(M_F) = (B_0^F | B_1'^{(F)} | \cdots | B_s'^{(F)}),$$

where $j \in \{1, \dots, m_F\}$, $B_j'^{(F)}$ is the matrix which columns are all the j-th columns of the blocks $B_1^{(F)}, \cdots, B_{m_F}^{(F)}$ ordered by the standard order. Let $\mathcal{F} = \{F \text{ facet of } \Gamma : \overline{F} \neq F\} = \{F_1, \dots, F_k\}$. We set $\Pi^* = (\Pi^*_{F_1}, \dots, \Pi^*_{F_k})$.

The next propositions is clear from the definition 4.14.

4.15 Proposition. *Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ and F a facet of Γ such that $F \neq \overline{F}$. Then Π^*_F is an admissible permutation (see definition 2.6) and consequently $\Pi^* \in \mathcal{P}_{\overline{\Gamma}}$.* ■

4.16 Proposition. *Let Γ be a clique complex and $\overline{\Gamma}$ a binomial extension of Γ. Let e be a virtual edge of G in $\overline{G}^{(\Pi^*)}$, such that $e = \{x_0^{(F)}, x_k^{(F)}\}$ for some facet F whose $Y_k^F \neq \emptyset$. Then:*

1. *If there exist $x \in (F \setminus V_\Gamma(M_F))$ such that for any vertex $x' \in V(C)$, $\{x, x'\} \notin E(\Gamma)$, then $t_\Pi(e) = 2$.*
2. *Else, let $J_F^{\text{ls}} = \{k\} \cup \{j \in \{1, \dots, m_F\} : \forall x' \in (V(C) \setminus e),\ \{x_j^{(F)}, x'\} \notin E(\Gamma)\}$:*
   - *(a) If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) < |Y_k^{(F)}|$, then $t_{\Pi^*}(e) = \min_{1 \leq j \leq m_F}(|Y_j^{(F)}|) + 2$.*
   - *(b) If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) = |Y_k^{(F)}|$, then $t_{\Pi^*}(e) = \min_{1 \leq j \leq m_F}(|Y_j^{(F)}|) + 1$.*

Proof. From the proposition 4.13 we have that the first statement is true. We should only prove the second statement

2. (a) By the definition of $\Pi^*_F(M_F)$ (see 2.10) we can describe all edges in $\overline{\Gamma}^{(\Pi^*)}$ inside $\overline{F}$, i.e.
   - $\{x_0, y_{j1}\}$ with $j \in \{1, \dots, m_F\}$; $\{x_0, x_1\}$ if $Y_1^{(F)} = \emptyset$.
   - For any $j, t \in \{1, \dots, m_F\}$, $i \in \{1, \dots, n_j\}$, $s \in \{1, \dots, n_t\}$ and $i \geq s$, $\{y_{ji}, y_{ts}\} \in \overline{\Gamma}^{(\Pi^*)}$ if and only if
     i. Either $i = s$ and $j \neq t$;
     ii. or $i = s + 1$ and $t \geq j$;
   - For any $j, t \in \{1, \dots, m_F\}$, $i \in \{1, \dots, n_j\}$ and $\varphi(t)$ the number in $\mathbb{N}$ such that x_t appears in the block $\varphi(t)$ of $\Pi^*(M_F)$; $\{y_{ji}, x_t\} \in \overline{\Gamma}^{(\Pi^*)}$ if and only if $i \geq \eta_e$ and
     i. either $\varphi(t) < i$;
     ii. or $\varphi(t) = i$ and $j \geq t$.
   - $\{x_j, x_{j'}\} \in E(\overline{\Gamma}^{(\Pi*)})$ for all $1 \leq j < j' \leq m_F$

   (b) We set $\eta_e = \min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|)$. Let $j' \in J_F^{\text{ls}}$ such that $|Y_{j'}^{(F)}| = \eta_e$, if $|Y_k^{(F)}| = \eta_e$, we set $j' = k$. We define P_e as
     i. if $j' \neq k$, $E(P_e) = \{\{x_0, y_{j'1}\}, \{y_{j'1}, y_{j'2}\}, \dots, \{y_{j'\eta_e}, x_{j'}\}, \{x_{j'}, x_k^{(F)}\}\}$; note that $|P_e| = \eta_e + 2$.
     ii. If $j' = k$, $E(P_e) = \{\{x_e, y_{k1}\}, \{y_{k1}, y_{k2}\}, \dots, \{y_{k\eta_e - 1}, y_{k\eta_e}\}, \{y_{kn_e}, x_k^{(F)}\}\}$; note that $|P_e| = \eta_e + 1$.

   By (a) P_e is a path in $\overline{\Gamma}^{(\Pi^*)}$ and does not have chords in F because Π^* is an admissible order. Moreover, since $j' \in J_F^{\text{ls}}$, there is not any edge from $V(P_e) \setminus e$ to a point $x \in V(C) \setminus e$. Thus P_e is a local substitution of e.

(c) We are going to prove that P_e is a local substitution of e of minimal lenght.
Let R be any local substitution of e. It is clear that R must start with an edge $\{x_0, y_{j_1,1}\}$ for $1 \leq j \leq m_F$ or $\{x_0, x_1\}$ if $1 \in J_F^{\text{ls}}$. In the case where R start with $\{x_0, x_1\}$, $\eta_e = 0$ and by (a) the stament is true. So we assume that R starts with $\{x_0, y_{j_1,1}\}$ for some $1 \leq j \leq m_F$. By (a) the last edge of R is:

- either $\{y_{ji}, x_k\}$ with $i \geq \eta_e$ and either $\varphi(k) < i$, or $\varphi(k) = i$ and $j \geq k$.
- Or $\{x_t, x_k\}$ with $t \in J_F^{\text{ls}}$. In this case there is an edge $\{y_{ji}, x_t\}$ with $i \geq \eta_e$ and either $\varphi(t) < i$ or $\varphi(t) = i$ and $j \geq t$. In any case there is $y_{ji} \in V(R)$, with $i \geq \eta_e$. Moreover, any other edge of R has the form $\{y_{ji}, y_{ts}\}$ with either $i = s$ and $j \neq t$; or $i = s+1$ and $t \geq j$. This prove that in order to go to y_{ji} with $i \geq \eta_e$ from $y_{j_1,1}$ we have to go forward or backward block by block, but we can never go forward more than one block by one step. Thus:
  i. If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) < |Y_k^{(F)}|$, then $|R| \geq i + 2 \geq \eta_e + 2 = |P_e|$.
  ii. If $\min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|) = |Y_k^{(F)}|$, then $|R| \geq i + 1 \geq \eta_e + 1 = |P_e|$.

So $t_{\Pi^*}(e) = P_e$. ■

4.17 Theorem. *Let* Γ *be a clique complex and* $\overline{\Gamma}$ *a binomial extension of* Γ *such that* $\mathcal{F} = \{F \text{ facet of } \Gamma : \overline{F} \neq F\}$ *is endowed by an admissible order. Then*

1. $p_2(\mathcal{B}_{\overline{\Gamma}}) \geq \min_{C \in \mathfrak{C}}(\min_{C' \in \mathfrak{C}_{\Pi^*}^{\text{ls}}(C)}|C'|) - 3.$

2. For any virtual edge $e \in (C)$ *and* F *the unique facet containing it, let* $J_F^{\text{ls}} = \{j \in \{1, \ldots, m_F\} : \forall x' \in (V(C) \setminus e),\ \{x_j^{(F)}, x'\} \notin E(\Gamma)\} \cup \{k\}$, $\eta_e = \min_{j \in J_F^{\text{ls}}}(|Y_j^{(F)}|)$,
$R^{(\Pi^*)}(C) = \{e \in E(C) | e \notin E(\overline{G}^{(\Pi^*)})\}$;
$R_1^{(\Pi^*)}(C) = \{e \in R^{(\Pi^*)}(C) | \exists x \in F \setminus V_\Gamma(M_F),\ \forall x' \in V(C),\ \{x, x\} \notin E(\overline{\Gamma})\}$;
$R_2^{(\Pi^*)}(C) = \{e \in R^{(\Pi^*)}(C) \setminus R_1^{(\Pi^*)}(C) | \exists k \in \{1, \ldots, m_F\},\ e \cap B_k^{(F)} \neq \emptyset, \text{ and } |Y_k^{(F)}| > \eta_e\}$;
$R_3^{(\Pi^*)}(C) = \{e \in R^{(\Pi^*)}(C) \setminus R_1^{(\Pi^*)}(C) | \exists k \in \{1, \ldots, m_F\},\ e \cap B_k^{(F)} \neq \emptyset, \text{ and } |Y_k^{(F)}| = \eta_e\}$. *Then, for any* $C \in \mathfrak{C}$ *and any* $C^{(\Pi^*)} \in \mathfrak{C}^{(\Pi^x)}(C)$,

$$C^{(\Pi^*)} = |C| + |R_1^{(\Pi^*)}(C)| + |R_2^{(\Pi^*)}(C)| + \sum_{e \in R_2^{(\Pi^*)}(C) \cup R_3^{(\Pi^*)}(C)} \eta_e.$$

Proof.

1. Thanks to proposition 4.10

$$p_2(\mathcal{B}_{\overline{\Gamma}}) \geq \max_{\Pi \in \mathcal{P}_{\overline{\Gamma}}}(\min_{C' \in \mathfrak{C}_\Pi^{\text{ls}}}|C'|) - 3,$$

Let $\Pi \in \mathcal{P}$. By definition of $\mathfrak{C}_\Pi^{\text{ls}}$ (c.f., 4.5) and proposition 4.16

$$\min_{C' \in \mathfrak{C}_{\Pi^*}^{\text{ls}}}|C'| = \min_{C \in \mathfrak{C}}(\min_{C' \in \mathfrak{C}_\Pi^{\text{ls}}(C)}|C'|).$$

And
$$\begin{aligned} \min_{C' \in \mathfrak{C}_\Pi^{\text{ls}}(C)}|C'| &= \min_{C \in \mathfrak{C}}(\textstyle\sum_{e \in E(C)} t_\Pi(e)) \\ &\leq \min_{C \in \mathfrak{C}}(\textstyle\sum_{e \in E(C)} t_{\Pi^*}(e)) \\ &= \min_{C' \in \mathfrak{C}_{\Pi^*}^{\text{ls}}(C)}|C'|. \end{aligned}$$

So $\max_{\Pi \in \mathcal{P}_{\overline{\Gamma}}}(\min_{C' \in \mathfrak{C}_\Pi^{\text{ls}}}|C'|) = \min_{C' \in \mathfrak{C}_{\Pi^*}^{\text{ls}}(C)}|C'|$ and we conclude that

$$p_2(\mathcal{B}_{\overline{\Gamma}}) \geq \min_{C \in \mathfrak{C}}(\min_{C' \in \mathfrak{C}_{\Pi^*}^{\text{ls}}(C)}|C'|) - 3.$$

2. Let $C \in \mathfrak{C}$ and $C^{(\Pi^*)} \in \mathfrak{C}^{(\Pi^x)}(C)$, using the proposition 4.16 we have that

$$\begin{aligned}
\textstyle\sum_{e\in C} t_{(\Pi^*)}(e) &= \textstyle\sum_{e\in E(C)-R^{(\Pi^*)}(C)} t_{\Pi^*}(e) + \sum_{e\in R_1^{(\Pi^*)}(C)} t(e) + \sum_{e\in R_2^{(\Pi^*)}(C)} t_{\Pi^*}(e) + \sum_{e\in R_3^{(\Pi^*)}(C)} t_{\Pi^*}(e)\\
&= \textstyle\sum_{e\in E(C)-R^{(\Pi^*)}(C)} 1 + \sum_{e\in R_1^{(\Pi^*)}(C)} 2 + \sum_{e\in R_2^{(\Pi^*)}(C)} (\eta_e+2) + \sum_{e\in R_3^{(\Pi^*)}(C)} (\eta_e+1)\\
&= \textstyle\sum_{e\in E(C)} 1 + \sum_{e\in R_1^{(\Pi^*)}(C)} 1 + \sum_{e\in R_2^{(\Pi^*)}(C)} 1 + \sum_{e\in R_2^{(\Pi^*)}(C)\cup R_3^{(\Pi^*)}(C)} (\eta_e)\\
&= |C| + |R_1(C)| + |R_2(C)| + \textstyle\sum_{e\in R_2^{(\Pi^*)}(C)\cup R_3^{(\Pi^*)}(C)} (\eta_e).
\end{aligned}$$

■

5 Upper bounds of $p_2(\mathcal{B}_{\overline{\Gamma}})$

5.1 Proposition. *Let C_n be the cycle of n-vertices $x_1,\ldots,x_n$ and Γ the clique complex generated by C_n. For each edge $\{x_k,x_{k+1}\}$ where $k=1,\ldots,n$ $(x_{n+1}=x_1)$ we associate a set Y_k of cardinality $s_i\in\mathbb{N}$ and we set $s=\sum_{j=1}^{m} s_i$. Then $\beta_{i,j}(\mathcal{B}_{\overline{\Gamma}}) = \beta_{i,j}(I_{\overline{\Gamma}'}) = (d+s)\frac{i}{d+s-i-1}\binom{d+s-2}{i+1}$ and $p_2(\mathcal{B}_{\overline{\Gamma}}) = p_2(I_{\overline{\Gamma}'}) = n+s-3$. In particular $\Bbbk[\mathbf{x}]/\mathcal{B}_\Gamma$ is Gorenstein.*

Proof. By remark 3.2, $\mathcal{F} = \{F \text{ facet of } C_n : \overline{F} \neq F\}$ is endowed by an admissible ordered. So, by theorem 2.12 $\mathrm{in}(\mathcal{B}_{\overline{\Gamma}}) = \mathrm{I}_{\overline{\Gamma}'}$. By definition 2.10 we have that $\overline{\Gamma}'$ is a $n+s$ cycle. By Hochster's formula we know that the free minimal of resolution $\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'}$ has the following shape:

$$0 \to \Bbbk[\mathbf{x}](-n-s) \to \Bbbk[\mathbf{x}]^{\beta_{n+s-3}}(-n-s+2) \to \cdots \to \Bbbk[\mathbf{x}]^{\beta_1}(-2) \to \Bbbk[\mathbf{x}] \to \Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'} \to 0,$$

where for $i=1,\ldots,n+s-3$,

$$\beta_i = \beta_{i,i+1}(\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'}) = \beta_{i-1,i+1}(I_{\overline{\Gamma}'}) \text{ and } 1 = \beta_{n+s-2,n+s}(\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'}) = \beta_{n+s-3,n+s}(I_{\overline{\Gamma}'}).$$

Then $p = p_2(I_{\overline{\Gamma}'}) = n+s-3$ and $\mathrm{dimproj}(\Bbbk[\mathbf{x}]/I) = n+s-2$, thus $\Bbbk[\mathbf{x}]/I$ is Cohen-Macaulay. By the proposition 1.4 and by the theorem 1.1 we deduce that

$$\beta_{i,j}(S/\mathcal{B}_{\overline{\Gamma}}) = \begin{cases} 0 & \text{si } j \neq i+1 \text{ and } i \neq n+s-2,\\ 0 & \text{si } i = n+s-2 \text{ and} j \neq n+s,\\ \beta_{i,j}(\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'}) & \text{si } j = i+1. \end{cases}$$

Thus projdim $\Bbbk[\mathbf{x}]/\mathcal{B}_{\bar{\Gamma}} \leq n+s-2$. Moreover, $\mathcal{K}(\Bbbk[\mathbf{x}]/\mathcal{B}_{\bar{\Gamma}};t) = \mathcal{K}(\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'};t)$. So the leading terms are equal and:

$$1 = \beta_{n+s-2,n+s}(\Bbbk[\mathbf{x}]/I_{\overline{\Gamma}'}) = \sum_{j=0}^{n+s-2} (-1)^{n+s-2-j}\beta_{j,n+s} = \beta_{n+s-2,n+s}.$$

So the free minimal of resolution of $\Bbbk[\mathbf{x}]/\mathcal{B}_\Gamma$ is:

$$0 \to \Bbbk[\mathbf{x}](-n-s) \to \Bbbk[\mathbf{x}]^{\beta_{n+s-3}}(-n-s+2)\ldots\Bbbk[\mathbf{x}]^{\beta_1}(-2) \to \Bbbk[\mathbf{x}] \to \Bbbk[\mathbf{x}]/B_{\overline{\Gamma}} \to 0.$$

As $\Bbbk[\mathbf{x}]/I_{\Gamma'}$ is Cohen-Macaulay and d-pure with $d=(0,2,3,\ldots,-2,n+s-2,n+s)$, by the Herzog-Kühl equations [12]

$$\beta_i = \beta_0(-1)^{i+1}\prod_{k\neq i,k\neq 0}\frac{d_k-d_0}{d_k-d_i},$$

for $1 \leq i \leq d+s-2 = -2$. We set $n := d+s$ and we have that:

$$\begin{aligned}\beta_{i-1}(\mathcal{B}_{\overline{\Gamma}}) &= \beta_{i-1}(\mathcal{I}_{\overline{\Gamma}}) = \beta_i = (-1)^{i+1}(\textstyle\prod_{k=1}^{i-1} \frac{k+1}{k+1-(i+1)})(\prod_{k=i+1}^{n-3} \frac{k+1}{k+1-(i+1)})(\frac{n-2}{n-2-(i+1)})\\ &= (-1)^{i+1}(-1)^{i-1}(\textstyle\prod_{k=1}^{i-1} \frac{k+1}{i-k})(\prod_{k=i+1}^{n-3} \frac{k+1}{k-i})(\frac{n}{n-(i+1)})\\ &= \tfrac{(n-2)!/i+1}{(i-1)!(n-3-i)!} \tfrac{n}{n-(i+1)} = n\tfrac{i}{n-i-1} \tfrac{(n-2)!}{\frac{(i+1)!}{i}(n-2-(i+1))!}\\ &= n\tfrac{i}{n-i-1} \tfrac{(n-2)!}{(i+1)!(n-2-(i+1))!} = n\tfrac{i}{n-i-1} \tbinom{n-2}{i+1}.\end{aligned}$$

In particular, we deduce that $p_2(\mathcal{B}_{\overline{\Gamma}}) = n + s - 3 = p_2(I_{\overline{\Gamma}'})$ and $\Bbbk[\mathbf{x}]/B_{\overline{\Gamma}}$ is Gorenstein. ■

Before proving our second principal result (cf. 5.9) we are going to introduce an special graduation for $\Bbbk[\mathbf{x}]$. By proposition 1.11 (cf. [7]) if B is a binomial ideal not including monomials, so the algebra $\Bbbk[\mathbf{x}]$ can be graded by a semigroup $\Sigma(B)$ in the following way: let $B \subset \Bbbk[\mathbf{x}]$ a binomial ideal not including monomials, so $\Sigma(B) = \mathbb{N}^n / \cong_B$ where

$$\forall \alpha, \beta \in \mathbb{N}^n,\ \alpha \cong_B \beta, \text{ if there exist a binomial } \mathbf{x}^\alpha - a\mathbf{x}^\beta \in B \text{ with } a \in \Bbbk.$$

We set the morphism between semigroups $\deg_{\Sigma_B} : \mathbb{N}^n \to \Sigma(B)$ defined by $\deg_{\Sigma(B)}(\alpha) = [\alpha]_{\cong B}$. So $\Bbbk[\mathbf{x}] = \oplus_{\sigma\in\Sigma(B)} S_\sigma$, where $S_\sigma = \oplus_{\deg(\alpha)=\sigma} \Bbbk \cdot x^\alpha$, is a $\Sigma(B)$-graduation of $\Bbbk[\mathbf{x}]$. If B is an toric homogeneous ideal of $\Bbbk[\mathbf{x}]$, we have that $\Bbbk[\mathbf{x}]$ is positively $\Sigma(B)$-graded.

5.2 Proposition. *[2, Proposition 1.1] Let B be a toric ideal and I be a squarefree monomial ideal of $\Bbbk[\mathbf{x}]$ such that: for any binomial $\mathbf{x}^\alpha - \mathbf{x}^\gamma \in B$, we have that $\mathbf{x}^\alpha \in I$ if and only if $x^\gamma \in I$. Let $J = B + I$; $R = \Bbbk[\mathbf{x}]/J$. Setting*

$$\begin{gathered}\Omega_b = \{F \subset \{1, \ldots, n\} : \exists \beta, \gamma \in \mathbb{N}^n,\ \deg_{\Sigma(B)}(\beta) = b,\ \mathbf{x}^\beta - \mathbf{x}^\gamma x^F \in B\} \text{ and}\\ \Psi_b = \{F \in \Omega_b : \exists \beta, \gamma \in \mathbb{N}^n,\ \deg_{\Sigma(B)}(\beta) = b,\ \mathbf{x}^\beta - \mathbf{x}^\gamma x^F \in B,\ \mathbf{x}^\gamma \in I\}.\end{gathered}$$

Then for all $b \in \Sigma(B)$ $\beta_{i,b}(R) = \dim_{\Bbbk} \widetilde{\mathrm{H}}_{i-1}(\Omega_b, \Psi_b(B); \Bbbk)$.

Now, we return to our problem. Let Γ be a clique complex with n vertices, $\overline{\Gamma}$ a binomial extension of Γ, $m = |V(\overline{\Gamma})|$. Moreover we will denote by $\mathbf{x} = (x_1, \ldots, x_n)$ the variable set identifying the vertex set of Γ, $\mathbf{y}$ the variable set identifying the vertices in $V(\overline{\Gamma}) \setminus V(\Gamma)$ and $\mathbf{z} = (\mathbf{x}, \mathbf{y})$. So I_Γ is an ideal of $\Bbbk[\mathbf{x}]$ and $\mathcal{B}_{\overline{\Gamma}}$ is an ideal of $\Bbbk[\mathbf{z}]$. We recall: let F be a facet of Γ we set $V_\Gamma(M_F)$ the set of all the vertices of Γ staying in M_F, if M_F is defined; otherwise $V_\Gamma(M_F) = \emptyset$. Besides $V_{\overline{\Gamma}}(M_F)$ are all the elements of $\overline{F}$ that appear in M_F.

5.3 Definition. Let $\mathbf{z_1}, \ldots, \mathbf{z_k}$ be variable sets and $I_1 \subset \Bbbk[\mathbf{z_1}], \ldots, I_k \subset \Bbbk[\mathbf{z_k}]$ be toric ideals such that for all pair $i, j \in \{1, \ldots, k\}$, $i \neq j$, $|\mathbf{z_i} \cap \mathbf{z_j}| \leq 1$. We define the **graph of the sequence of toric ideals** $I_1, \ldots, I_k$, denoted by $G(I_1, \ldots, I_k)$,as the graph whose vertice set is $I_1, \ldots, I_k$ and $\{I_i, I_j\}$ is an edge if I_i and I_j has a common variable.

5.4 Theorem. ([4]) *Let $\mathbf{z_1}, \ldots, \mathbf{z_k}$ be variable sets and $I_1 \subset \Bbbk[\mathbf{z_1}], \ldots, I_k \subset \Bbbk[\mathbf{z_k}]$ be toric ideals such that for any pair $i, j \in \{1, \ldots, k\}$, $i \neq j$, $|\mathbf{z_i} \cap \mathbf{z_j}| \leq 1$. If for any connex component $G(I_1, \ldots, I_k)$ is a tree, then $J := \sum_{i=1}^k I_i$ is a homogeneous toric ideal and setting r as the number of connexes components of $G(I_1, \ldots, I_k)$ we have that:*

$$\dim\ (\Bbbk[\mathbf{x}]/J) = \sum_{i=1}^{k} \dim\ (\Bbbk[\mathbf{z_i}]/I_i) + r - k + 1,$$

where $\mathbf{x} = \cup_{i=1}^k \mathbf{z_i}$

5.5 Lemma. *Let Γ be a clique complex, $\overline{\Gamma}$ a binomial extension of Γ such that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. Let $F \in \mathcal{F} = \{F \text{ facet of } \Gamma : \overline{F} \neq F\}$. For all 2×2 minor of M_F, $\mathbf{z}^\alpha - \mathbf{z}^\beta$, and all $\gamma \in \mathbb{N}^m$ we have that $\mathbf{z}^{\gamma+\alpha} \in \mathcal{I}_{\overline{\Gamma}}$ if and only if $\mathbf{z}^{\gamma+\beta} \in \mathcal{I}_{\overline{\Gamma}}$. In particular if $\mathbf{z}^{\gamma+\alpha} \in \mathcal{I}_{\overline{\Gamma}}$, then $\mathbf{z}^\gamma(\mathbf{z}^\alpha - \mathbf{z}^\beta) \in \mathcal{I}_{\overline{\Gamma}}$.*

Proof. To show that $\mathbf{z}^{\gamma+\alpha} \in \mathcal{I}_{\overline{\Gamma}}$ if and only if $\mathbf{z}^{\gamma+\beta} \in \mathcal{I}_{\overline{\Gamma}}$, it is enough to show that if $\mathbf{z}^{\gamma+\alpha} \in \mathcal{I}_{\overline{\Gamma}}$ then $\mathbf{z}^{\gamma+\beta} \in \mathcal{I}_{\overline{\Gamma}}$, since $\mathbf{z}^\beta - \mathbf{z}^\alpha = -(\mathbf{z}^\alpha - \mathbf{z}^\beta)$ is also a 2 minor of M_F.

As $\mathbf{z}^\alpha - \mathbf{z}^\beta$ is a 2×2 minor of M_F, γ has to be different to 0, otherwise $\mathbf{z}^\alpha \in I_\Gamma$, but by definition of M_F, is not possible. Furthermore, as $\mathbf{z}^{\gamma+\alpha} \in \mathcal{I}_{\overline{\Gamma}}$, there exist $z_1, z_2 \in V(\overline{\Gamma})$ such that $z_1 z_2 \in \mathcal{I}_{\overline{\Gamma}}$ and $z_1 z_2 | z^{\gamma+\alpha}$. We have three cases:

1. Let $z_1 | \mathbf{z}^\alpha$ and $z_2 | \mathbf{z}^\alpha$. Thus $z^\alpha \in I_\Gamma$, but we have already said that by definition of M_F, it is not possible.

2. Let $z_1 | \mathbf{z}^\alpha$ or $z_2 | \mathbf{z}^\alpha$ but no both in the same time. Then, we can suppose $z_1 | \mathbf{z}^\alpha$ and $z_2 \nmid \mathbf{z}^\alpha$; so $z_2 | \mathbf{z}^\gamma$. As $z_1 | \mathbf{z}^\alpha$, we have that $z_1 \in \overline{F}$. Moreover $z_1 z_2 \in \mathcal{I}_{\overline{\Gamma}}$, thus $z_2 \notin \overline{F}$ and for all F' facet of Γ such that $z_2 \in \overline{F'}$ $z_1 \notin \overline{F'}$. As $\mathbf{z}^\alpha - \mathbf{z}^\beta$ is a minor 2×2 of M_F, there exist $z_1', z_2' \in V_{\overline{\Gamma}}(M_F) \subset \overline{F}$ such that $\mathbf{z}^\beta = z_1' z_2'$ and we have two cases:
   (a) $z_1' = z_2'$. Then $z_1' \in Y^{(F)}$ and z_1' is only in $\overline{F}$.

   (b) $z_1' \neq z_2'$. Then $z_1' \notin \overline{F}'$ or $z_2' \notin \overline{F}'$ for all F' facet de Γ such that $z_2 \in \overline{F}'$. Otherwise, if there exist a facet F' of Γ such that $z_2 \in \overline{F}'$ and $z_1', z_2' \in \overline{F}'$, then $\{z_1', z_2'\} \in \overline{F} \cap \overline{F'}$, so $\{z_1', z_2'\} \in F \cap F'$ and $\{z_1', z_2'\}$ is not a proper edge of Γ. As $\mathbf{z}^\alpha - z_1' z_2'$ is a 2×2 minor of M_F, we have that $x_0^F \in \{z_1', z_2'\}$, thus, by definition of $\mathcal{B}_{\overline{\Gamma}}$, $\{z_1', z_2'\}$ is an edge proper of Γ. Consequently $z_1' \notin \overline{F'}$ or $z_2' \notin \overline{F'}$ for all F' facet of Γ such that $z_2 \in \overline{F}'$.

   In this way, for the two cases ther exist $z' \in \overline{F}$ such that $z' | \mathbf{z}^\beta$ and $z' \notin \overline{F'}$ for all F' facet of Γ such that $z_2 \in \overline{F'}$. Consequently $z' z_2 \in I_\Gamma$. Moreover $z' z_2 | \mathbf{z}^{\gamma+\beta}$, so $\mathbf{z}^{\gamma+\beta} \in \mathcal{I}_{\overline{\Gamma}}$.

3. Let $z_1 \nmid \mathbf{z}^\alpha$ and $z_2 \nmid \mathbf{z}^\alpha$. Then $z_1 z_2 | \mathbf{z}^\gamma$. So $z_1 z_2 | \mathbf{z}^{\gamma+\beta}$ an in this way $\mathbf{z}^{\gamma+\beta} \in \mathcal{I}_{\overline{\Gamma}}$.

■

5.6 Proposition. *Let be G a graph, $\Gamma = \Gamma(G)$ its clique complex, $\overline{\Gamma}$ a binomial extension of Γ such that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. If $\mathbf{z}^\alpha - \mathbf{z}^\beta \in \mathcal{J}_{\overline{\Gamma}}$ such that $\mathbf{z}^\alpha \in \mathcal{I}_{\overline{\Gamma}}$, then $\mathbf{z}^\beta \in \mathcal{I}_{\overline{\Gamma}}$.*

Proof. As $\mathbf{z}^\alpha - \mathbf{z}^\beta \in \mathcal{J}_{\overline{\Gamma}}$, there exist $a_1, \ldots, a_n \in \Bbbk$, $\gamma_1, \ldots, \gamma_n \in \mathbb{N}^m$, $F_1, \ldots, F_n$ facets of Γ and $b_1, \ldots, b_n$ 2 minors of $M_{F_1}, \ldots, M_{F_n}$ respectively, such that $\mathbf{z}^\alpha - \mathbf{z}^\beta = \sum_{i=1}^n a_i \mathbf{z}^{\gamma_i} b_i$. Let

$$\mathcal{K} := \{i \in \{1, \ldots, n : \text{ one of the monomials of } z^{\gamma_i} b_i \in \mathcal{I}_{\overline{\Gamma}}\}.$$

$\mathcal{K} \neq \emptyset$, since there must exist $i \in \{1, \ldots, n\}$ such that z^α is a monomial of $\mathbf{z}^{\gamma_i} b_i$. By the lemma 5.5, if one of the monomials of $\mathbf{z}^{\gamma_i} b_i \in \mathcal{I}_{\overline{\Gamma}}$, then the two monomials of $z^{\gamma_i} b_i$ belong to $\mathcal{I}_{\overline{\Gamma}}$. So

$$z^\alpha - z^\beta = \sum_{i \in \mathcal{K}} a_i \mathbf{z}^{\gamma_i} b_i + \sum_{i \notin \mathcal{K}} a_i \mathbf{z}^{\gamma_i} b_i.$$

- If z^β appears in a term of $\sum_{i \in \mathcal{K}} a_i \mathbf{z}^{\gamma_i} b_i$, then $z^\beta \in I_\Gamma$.

- If z^{β} does not appear as a term of $\sum_{i\in\mathcal{K}} a_i\mathbf{z}^{\gamma_i}b_i$, then z^{β} must to appear as a term of $\sum_{i\notin\mathcal{K}} a_i\mathbf{z}^{\gamma_i}b_i$. Since all monomial of the first addition does not appear as a term of the second addition, we have:

  $$z^{\alpha}-\sum_{i\in\mathcal{K}} a_i\mathbf{z}^{\gamma_i}b_i = z^{\beta}-\sum_{i\notin\mathcal{K}} a_i\mathbf{z}^{\gamma_i}b_i = 0.$$

  Thus $z^{\alpha}\in\mathcal{J}_{\overline{\Gamma}}$, but it is a contradiction to the fact that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. So this case is not possible.

$\blacksquare$

5.7 Lemma. *Let G be a graph, $\Gamma=\Gamma(G)$ be its clique complex, $\overline{\Gamma}$ be a binomial extension of Γ with $m=|V(\overline{\Gamma})|$, such that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. Let C be a virtual minimal cycle of G satisfying that for any edge e virtual, $e=\{x_0^{(F_e)},x_1^{(F_e)}\}$, where F_e is the only facet of Γ containing e. Let $\widetilde{C}$ be the cycle obtained from C replacing any virtual edge e by the edges $\{x_0^{(F_e)},y_{11}^{(F_e)}\},\{y_{11}^{(F_e)},y_{12}^{(F_e)}\},\ldots,\{y_{1n_1}^{(F_e)},x_1^{(F_e)}\}$. Then:*

$$\beta_{|\widetilde{C}|-3,\deg_{\Sigma_{\mathcal{J}_{\overline{\Gamma}}}}(z^{V(\widetilde{C})})}(\mathcal{B}_{\overline{\Gamma}})\geq 1.$$

Proof. We set $b=\deg_{\mathcal{J}_{\overline{\Gamma}}}\mathbf{z}^{V(\widetilde{C})}$. by the proposition 5.6, if $\mathbf{z}^{\gamma}-\mathbf{z}^{\beta}\in\mathcal{J}_{\overline{\Gamma}}$ and $\mathbf{z}^{\gamma}\in\mathcal{I}_{\overline{\Gamma}}$, then $\mathbf{z}^{\beta}\in\mathcal{I}_{\overline{\Gamma}}$. In this way we may use the Bruns-Herzog formula (cf. 5.2) to calculate $\beta_{|V(\widetilde{C})|-3,b}$:

$$\beta_{|\widetilde{C}|-3,b}(\mathcal{B}_{\overline{\Gamma}})=\beta_{|\widetilde{C}|-2,b}(\Bbbk[\mathbf{z}]/\mathcal{B}_{\overline{\Gamma}})=\dim_{\Bbbk}\widetilde{\mathrm{H}}_{|V(\widetilde{C})|-3}(\Omega_b,\Psi_b;\Bbbk),$$

where

$$\Omega_b=\{F\subset V(\overline{\Gamma}):\exists\beta,\gamma\in\mathbb{N}^m,\ \deg_{\Sigma(\mathcal{J}_{\overline{\Gamma}})}(\beta)=b,\ \mathbf{z}^{\beta}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)}\in\mathcal{J}_{\overline{\Gamma}}\}\text{ and}$$
$$\Psi_b=\{F\in\Omega_b:\exists\beta,\gamma\in\mathbb{N}^m,\ \deg_{\Sigma(\mathcal{J}_{\overline{\Gamma}})}(\beta)=b,\ \mathbf{z}^{\beta}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)}\in\mathcal{J}_{\overline{\Gamma}},\ \mathbf{z}^{\gamma}\in\mathcal{I}_{\overline{\Gamma}}\}.$$

- Firstly we show that: $\Omega_b=\{F\subset V(\overline{\Gamma}):\exists\gamma\in\mathbb{N}^m,\ \mathbf{z}^{\alpha}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)}\in\mathcal{J}_{\overline{\Gamma}}\}$.

  $\deg_{\Sigma(\mathcal{J}_{\overline{\Gamma}})}(\beta)=b$ if and only if $(\mathbf{z}^{\alpha}-\mathbf{z}^{\beta})\in\mathcal{J}_{\overline{\Gamma}}$. Then $\mathbf{z}^{\beta}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)}\in\mathcal{J}_{\overline{\Gamma}}$ if and only if $(\mathbf{z}^{\alpha}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)})\in\mathcal{J}_{\overline{\Gamma}}$. Consequently $\Omega_b=\{F\subset V(\overline{\Gamma}):\exists\gamma\in\mathbb{N}^m,\ \mathbf{z}^{\alpha}-\mathbf{z}^{\gamma}\mathbf{z}^{\mathrm{supp}(F)}\in\mathcal{J}_{\overline{\Gamma}}\}$.

- We are going to show that for any facet F of Γ such that $M_F\neq 0$ we have that any 2×2 minor of M_F $b(\mathbf{z})=z_{i,1}z_{i,2}-z_{i,3}z_{i,4}$ satisfies that: $z_{i,1},z_{i,2}\in V(\widetilde{C})$ if and only if $z_{i,3},z_{i,4}\in V(\widetilde{C})$.
  Firstly we recall the definition of M_F:

  $$M_F=\left(\begin{array}{cccc|ccc|c|ccc} x_0^{(F)} & y_{11}^F & \cdots & y_{1n_1}^F & y_{21}^F & \cdots & y_{2n_2}^F & \cdots & y_{m_F1}^F & \cdots & y_{m_Fn_{m_F}}^F \\ y_{11}^F & y_{12}^F & \cdots & x_1^F & y_{22}^F & \cdots & x_2^F & \cdots & y_{m_F2}^F & \cdots & x_{m_F}^F \end{array}\right);$$

  and

  $$\mathcal{I}_F=\{[M_F]_{1,u}[M_F]_{2,v}-[M_F]_{1,v}[M_F]_{2,u}:1\leq u<v\leq(\sum_{j=1}^{m_F}n_j)+1\}.$$

  As the restiction of $\widetilde{C}$ in F is $\{x_0^{(F)},y_{11}^{(F)}\},\{y_{11}^{(F)},y_{12}^{(F)}\},\ldots,\{y_{1n_1}^{(F)},x_1^{(F)}\}$, any minor $b(z)$ of M_F has all its variables in $V(\widetilde{C})$ if and only if $b(z)$ is a minor of the first block of M_F; so the second monomial $b(z)$ have also all its variables in $V(\widetilde{C})$. So we conclude the assertion.

- We are going to show that $\Omega_b =< V(\widetilde{C}) >$ (the simplicial complex whose all faces are all the subsets of $V(\widetilde{C})$).

  Let $\mathbf{z}^\alpha - \mathbf{z}^\beta \in \mathcal{J}_{\overline{\Gamma}}$. We suppose that some variable of $\mathbf{z}^\beta$ is not in $V(\widetilde{C})$. As $\mathbf{z}^\alpha - \mathbf{z}^\beta \in \mathcal{J}_{\overline{\Gamma}}$, there exist $a_1, \dots, a_n \in \Bbbk$, $\gamma_1, \dots, \gamma_n \in \mathbb{N}^m$, $F_{s_1}, \dots, F_{s_n}$ facets of Γ and $b_1 = \mathbf{z}^{\alpha_1} - \mathbf{z}^{\beta_1}, \dots, b_n = \mathbf{z}^{\alpha_n} - \mathbf{z}^{\beta_n}$ 2×2 minors of the matrices $M_{F_{s_1}}, \dots, M_{F_{s_n}}$ respectively, such that $\mathbf{z}^\alpha - \mathbf{z}^\beta = \sum_{i=1}^n a_i \mathbf{z}^{\gamma_i} b_i$. Let

  $$\mathcal{K} := \{i \in \{1, \dots, n\} : \text{ a monomial of } \mathbf{z}^{\gamma_i} b_i \text{ has all its variables in } V(\widetilde{C})\}.$$

  $\mathcal{K} \neq \emptyset$, since there must exist $i \in \{1, \dots, n\}$ such that $\mathbf{z}^\alpha$ is a term of any $\mathbf{z}^{\gamma_i} b_i$. Moreover, we note that if a monomial of $\mathbf{z}^{\gamma_i} b_i$ has all its variables in $V(\widetilde{C})$, then all the variables of $\mathbf{z}^{\gamma_i}$ are in $V(\widetilde{C})$ and by the last statement, any variable of the two monomials of b_i are in $V(\widetilde{C})$. Thus, we can write $z^\alpha - z^\beta = \sum_{i \in K} a_i \mathbf{z}^{\gamma_i} b_i + \sum_{i \notin K} a_i \mathbf{z}^{\gamma_i} b_i$. And by this remark, if $i \notin K$, any monomial of $\mathbf{z}^{\gamma_i} b_i$ does not have all its variables in $V(\widetilde{C})$, so:

  $$z^\alpha + \sum_{i \in K} a_i \mathbf{z}^{\gamma_i} b_i = z^\beta + \sum_{i \notin K} a_i \mathbf{z}^{\gamma_i} b_i = 0.$$

  Thus $z^\alpha \in \mathcal{J}_{\overline{\Gamma}}$, that is a contradiction to the fact that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. Then all the variables of $\mathbf{z}^\beta$ are in $V(\widetilde{C})$. And in this we way we can conclude:

  $$\begin{aligned} \Omega_b &= \{F \subset V(\overline{\Gamma}) : \exists \gamma \in \mathbb{N}^m,\ \mathbf{z}^\alpha - \mathbf{z}^\gamma \mathbf{z}^F \in \mathcal{J}_{\overline{\Gamma}}\} \\ &= \{F \subset V(\widetilde{C}) : \exists \gamma \in \mathbb{N}^m,\ \mathbf{z}^\alpha - \mathbf{z}^{\gamma + \mathrm{supp}(F)} \in \mathcal{J}_{\overline{\Gamma}}\} \\ &=< V(\widetilde{C}) > . \end{aligned}$$

- Let us show that $\Psi_b = (\overline{\Gamma}_{V(\widetilde{C})})^A$.
  By the same way that we showed that $\Omega_b = \{F \subset V(\overline{\Gamma}) : \exists \gamma \in \mathbb{N}^s,\ \mathbf{z}^\alpha - \mathbf{z}^\gamma \mathbf{z}^{\mathrm{supp}(F)} \in \mathcal{J}_{\overline{\Gamma}}\}$, we can prove that

  $$\Psi_b = \{F \in \Omega_b : \exists \gamma \in \mathbb{N}^m,\ \mathbf{z}^\alpha - \mathbf{z}^\gamma \mathbf{z}^{\mathrm{supp}(F)} \in \mathcal{J}_{\overline{\Gamma}} \text{ and } \mathbf{z}^\gamma \in \mathcal{I}_{\overline{\Gamma}}\}.$$

  As $\Omega_b =< V(\widetilde{C}) >$, we have that

  $$\begin{aligned} \Psi_b &= \{F \in V(\widetilde{C}) : \exists \gamma \in \mathbb{N}^m,\ \mathbf{z}^{\mathrm{supp}(F) + \mathrm{supp}(V(\widetilde{C}) \setminus F)} - \mathbf{z}^\gamma \mathbf{z}^{\mathrm{supp}(F)} \in \mathcal{J}_{\overline{\Gamma}} \text{ and } \mathbf{z}^\gamma \in \mathcal{I}_{\overline{\Gamma}}\} \\ &= \{F \in V(\widetilde{C}) : \exists \gamma \in \mathbb{N}^m,\ \mathbf{z}^{\mathrm{supp}(F)}(\mathbf{z}^{\mathrm{supp}(V(\widetilde{C}) \setminus F)} - \mathbf{z}^\gamma) \in \mathcal{J}_{\overline{\Gamma}} \text{ and } \mathbf{z}^\gamma \in \mathcal{I}_{\overline{\Gamma}}\}. \end{aligned}$$

  But $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal, so

  $$\mathcal{J}_{\overline{\Gamma}} = (\mathcal{J}_{\overline{\Gamma}} : (z_1 \cdots z_m)^\infty) = \{p(\mathbf{z}) \in \Bbbk[z_1 \cdots z_m] : \exists \alpha \in \mathbb{N}^m,\ \mathbf{z}^\alpha p(z) \in \mathcal{J}_{\overline{\Gamma}}\},$$

  then

  $$\Psi_b = \{F \in V(\widetilde{C}) : \exists \gamma \in \mathbb{N}^m,\ (\mathbf{z}^{\mathrm{supp}(V(\widetilde{C}) \setminus F)} - \mathbf{z}^\gamma) \in \mathcal{J}_{\overline{\Gamma}} \text{ and } \mathbf{z}^\gamma \in \mathcal{I}_{\overline{\Gamma}}\}.$$

  Besides, by the proposition 5.6 we deduce:

  $$\begin{aligned} \Psi_b &= \{F \in V(\widetilde{C}) : \exists \gamma \in \mathbb{N}^m,\ (\mathbf{z}^{\mathrm{supp}(V(\widetilde{C}) \setminus F)} - \mathbf{z}^\gamma) \in \mathcal{J}_{\overline{\Gamma}} \text{ and } \mathbf{z}^\gamma, \mathbf{z}^{\mathrm{supp}(V(\widetilde{C}) \setminus F)} \in \mathcal{I}_{\overline{\Gamma}}\} \\ &= \{F \in V(\widetilde{C}) : \mathbf{z}^{\mathrm{supp}(V(\widetilde{C}) \setminus F)} \in \mathcal{I}_{\overline{\Gamma}}\} \\ &= \{F \in V(\widetilde{C}) : (V(\widetilde{C}) \setminus F) \notin \overline{\Gamma}\} \\ &= \{F \in V(\widetilde{C}) : (V(\widetilde{C}) \setminus F) \notin \overline{\Gamma}_{V(\widetilde{C})}\} \\ &= (\overline{\Gamma}_{V(\widetilde{C})})^A. \end{aligned}$$

Using the Alexander duality and the long exact sequence of relative homology we have that

$$\begin{aligned}
\beta_{|\widetilde{C}|-3,b}(\mathcal{B}_{\overline{\Gamma}}) &= \dim_{\Bbbk}\widetilde{\mathrm{H}}_{|\widetilde{C}|-3}(\Omega_b,\Psi_b;\Bbbk) \\
&= \dim_{\Bbbk}\,\widetilde{\mathrm{H}}_{|\widetilde{C}|-(|\widetilde{C}|-3)-2}((\Psi_b)^A,(\Omega_b)^A;\Bbbk) \\
&= \dim_{\Bbbk}\,\widetilde{\mathrm{H}}_1((\Psi_b)^A;\Bbbk) \\
&= \dim_{\Bbbk}\,\widetilde{\mathrm{H}}_1(\overline{\Gamma}_{V(\widetilde{C})};\Bbbk).
\end{aligned}$$

- We are going to prove that $\dim_{\Bbbk}\,\widetilde{\mathrm{H}}_1(\overline{\Gamma}_{V(\widetilde{C})};\Bbbk)\geq 1$.

  We note that the facets of $\overline{\Gamma}_{V(\widetilde{C})}$ are the subset $\widetilde{F}_e$ for all $e\in E(C)$ defined by:

$$\widetilde{F}_e=\begin{cases} e & \text{if } e\in E(\widetilde{C}),\\ e\cup Y_1^{(F_e)} & \text{if } e\notin E(\widetilde{C}).\end{cases}$$

  So $\overline{\Gamma}_{V(\widetilde{C})} =< \widetilde{F}_e : e\in C >$ and is homotopically equivalent to $\Gamma_{V(C)}$, as we can see in the figure 4, where we set $E(C)=\{\{x_1,x_2\},\{x_2,x_3\},\ldots,\{x_n,x_1\}\}$ and

$$\text{if } \{x_i,x_{i+1}\}\in E(\widetilde{C}),\ \widetilde{F}_{x_i,x_{i+1}}=\{x_i,x_{i+1},y^i_{11},\ldots,y^i_{1n_1}\},\text{ where } x_{n+1}=x_1.$$



$(\overline{\Gamma}_{V(\widetilde{C})})_1$ $(\overline{\Gamma}_C)_1$

Figure 4:

We know that $\widetilde{H}_1(\Gamma_{V(C)};\Bbbk)$ is the homology of the complex

$$\oplus_{\sigma\in(\Gamma_{V(C)})_{(2)}}\Bbbk\cdot f_\sigma \xrightarrow{\partial_2} \oplus_{\sigma\in E(\Gamma_{V(C)})}\Bbbk\cdot f_\sigma \xrightarrow{\partial_1} \oplus_{\sigma\in V(C)}\Bbbk\cdot f_\sigma,$$

where $\Gamma_{V(C)})_{(2)}=\{\sigma\in\Gamma_{V(C)} : |\sigma|=3\}$ and $\partial_i(e_\sigma)=\sum_{i\in\sigma}\mathrm{sign}(i,\sigma)e_{\sigma\setminus\{i\}}$, for any $i\in\{1,2\}$ and $\mathrm{sign}(i,\sigma)=(-1)^{r-1}$, if i is the r-th element of the set $\sigma\subset\{x_1<\cdots<x_n\}$, writen by increassing order.

We have that $\partial_1(\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}}-f_{\{x_i,x_{i+1}\}})=0$, so $\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}}-f_{\{x_1,x_n\}}\in\ker(\partial_1)$. Now we need to prove that there is not an element $x\in\oplus_{\sigma\in(\Gamma_{V(C)})_2}\Bbbk\cdot f_\sigma$ such that $\partial_2(x)=\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}}-f_{\{x_1,x_n\}}$. Otherwise, assume that there exist $x=\sum_{\sigma\in(\Gamma_{V(C)})_{(2)}} a_\sigma f_\sigma$ such that $\partial_2(x)=\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}}-f_{\{x_1,x_n\}}$.

We know that $\widetilde{H}_1(\Gamma_{V(C)};\Bbbk)$ is the homology of the complex

$$\oplus_{\sigma\in(\Gamma_{V(C)})_{(2)}}\Bbbk\cdot f_\sigma \xrightarrow{\partial_2} \oplus_{\sigma\in E(\Gamma_{V(C)})}\Bbbk\cdot f_\sigma \xrightarrow{\partial_1} \oplus_{\sigma\in V(C)}\Bbbk\cdot f_\sigma,$$

where $(\Gamma_{V(C)})_{(2)}=\{\sigma\in\Gamma_{V(C)} : |\sigma|=3\}$ and $\partial_i(e_\sigma)=\sum_{i\in\sigma}\mathrm{sign}(i,\sigma)e_{\sigma\setminus\{i\}}$, for any $i\in\{1,2\}$ and $\mathrm{sign}(i,\sigma)=(-1)^{r-1}$, if i is the r-th element of the set $\sigma\subset\{x_1<\cdots<x_n\}$, written by increasing

order.

We have that $\partial_1(\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}} - f_{\{x_i,x_{i+1}\}}) = 0$, so $\sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}} - f_{\{x_1,x_n\}} \in \ker(\partial_1)$. Now we need to prove that there is not an element $x \in \oplus_{\sigma\in(\Gamma_{V(C)})_2}\Bbbk \cdot f_\sigma$ such that $\partial_2(x) = \sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}} - f_{\{x_1,x_n\}}$. On the contrary, assume that there exist $x = \sum_{\sigma\in(\Gamma_{V(C)})_{(2)}} a_\sigma f_\sigma$ such that $\partial_2(x) = \sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}} - f_{\{x_1,x_n\}}$.

Let remark that since C is a virtual minimal cycle, any chord of C is not a chord in $\widetilde{C}$; then any chord $\{A, B\}$ of C, is a proper edge of C, and if F is the unique facet containing $\{A, B\}$, then either $A = [M_F]_{11}$ or $B = [M_F]_{11}$.

Let $e \in E(C)$, since e appears in $\partial_2(x)$ there exist $\sigma = \{x_{i_1}, x_{i_2}, x_{i_3}\} \in (\Gamma_{V(C)})_{(2)}$ such that $a_\sigma \neq 0$ and $\{x_{i_1}, x_{i_2}\} = e$. Neither $\{x_{i_2}, x_{i_3}\} \in E(C)$ or $\{x_{i_1}, x_{i_3}\} \in E(C)$, by item 2. of definition 4.1. Thus $\{x_{i_2}, x_{i_3}\}$ and $\{x_{i_1}, x_{i_3}\}$ are chords of C; since $\{x_{i_2}, x_{i_3}\}$ is in the image $\partial_2(\sigma)$ but doesn't appears in $\partial_2(x)$, then there exist $\sigma' = \{x_{i_2}, x_{i_3}, x_{i_4}\} \in (\Gamma_{V(C)})_{(2)}$ with $a'_\sigma \neq 0$. By item 1. of definition 4.1, $\{x_{i_2}, x_{i_3}\}$ is a chord of C, so it is a proper edge, and there exist a unique facet F of Γ such that $\sigma, \sigma' \subset F$ and $|F| \geq 4$. As $\{x_{i_2}, x_{i_3}\}$ and $\{x_{i_1}, x_{i_3}\}$ are chords of C, we should have $[M_F]_{11} = x_{i_3}$. By item 1. of definition 4.1, $\{x_{i_4}, x_{i_1}\}$ is a chord of C and thus either $x_{i_4} = [M_F]_{11}$ or $x_{i_1} = [M_F]_{11}$, but $[M_F]_{11} = x_{i_3}$. This is a contradiction. Thus, there is not an element $x \in \oplus_{\sigma\in(\Gamma_{V(C)})_2}\Bbbk \cdot f_\sigma$ such that $\partial_2(x) = \sum_{i=1}^{n-1} f_{\{x_i,x_{i+1}\}} - f_{\{x_1,x_n\}}$. So, $\dim_k \widetilde{H}_1(\Gamma_{V(C)}; \Bbbk) \geq 1$.

Thus $\beta_{|V(\widetilde{C})|-3,b}(\mathcal{B}_{\overline{\Gamma}}) = \dim_\Bbbk \widetilde{\mathrm{H}}_{|V(\widetilde{C})|-3}(\Omega_b, \Psi_b; \Bbbk) = \dim_\Bbbk \widetilde{\mathrm{H}}_1(\overline{\Gamma}_{V(\widetilde{C})}; \Bbbk) \geq 1$.

■

5.8 Definition. *We define* $\widetilde{\mathfrak{C}} = \{C \in \mathfrak{C}^v : \forall e \in E(C) \text{ virtual } e \in B_1(M_{F_e})\}$.

5.9 Theorem. *Let Γ be a clique complex and $\overline{\Gamma}$ a binomial extension of Γ such that $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. If $\widetilde{\mathfrak{C}} \neq \emptyset$, then $p_2(\mathcal{B}_{\overline{\Gamma}}) + 3 \leq \min_{C\in\widetilde{\mathfrak{C}}}(|\widetilde{C}|) = \min_{C\in\widetilde{\mathfrak{C}}}(|C| + \sum_{e\in R(C)} Y_1^{F_e})$.*

Proof. Let $C \in \widetilde{\mathfrak{C}}$. Consider $\widetilde{C}$ the cycle defined in lemma 5.7, so by this lemma:

$$\begin{aligned} \beta_{|\widetilde{C}|-3,|\widetilde{C}|}(\mathcal{B}_{\overline{\Gamma}}) &= \textstyle\sum_{h\in\Sigma(\mathcal{J}_{\overline{\Gamma}}),|h|=|\widetilde{C}|} \beta_{|\widetilde{C}|-3,h}(\mathcal{B}_{\overline{\Gamma}}) \\ &\geq \beta_{|\widetilde{C}|-3,\deg_{\Sigma_{\mathcal{J}_{\overline{\Gamma}}}}(z^{\mathrm{supp}(V(\widetilde{C}))})}(\mathcal{B}_{\overline{\Gamma}}) = 1, \end{aligned}$$

then $p_2(\mathcal{B}_{\overline{\Gamma}}) \leq |\widetilde{C}| - 3 = |C| + \sum_{e\in R(C)} Y_1^{(F_e)} - 3$. So

$$p_2(\mathcal{B}_{\overline{\Gamma}}) + 3 \leq \min_{C\in\widetilde{\mathfrak{C}}}(|\widetilde{C}|) = \min_{C\in\widetilde{\mathfrak{C}}}(|C| + \sum_{e\in R(C)} Y_1^{(F_e)}).$$

■

5.10 Theorem. *Let Γ be a clique complex and $\overline{\Gamma}$ be a binomial extension of Γ such that $\mathcal{F} = \{F$* facet of Γ : *$F \neq \overline{F}\}$ is endowed with an admissible order and $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. Suppose that $\widetilde{\mathfrak{C}} = \mathfrak{C}^v$, moreover for any $C \in \widetilde{\mathfrak{C}}$ and for any edge $e \in E(C)$ with F_e is the only facet that contains e, e satisfies one of the next properties:*

1. *either $|Y_1^{(F_e)}| = \min_{1\leq j\leq m_{F_e}}(Y_i^{(F_e)}) \geq 2$ and $V_\Gamma(M_{F_e}) = F_e$, où $J_{F_e}^{\mathrm{ls}} = \{j \in (\{1, \ldots, m_{F_e}\} \setminus \{k\}) : \forall x' \in (V(C) \setminus e),\ \{x_j^{(F)}\} \cup \{1\}$;*

2. or $|Y_1^{(F_e)}| = 1$.

Then, $p_2(\mathcal{B}_{\overline{\Gamma}}) = \min_{C\in\mathfrak{C}^v}(|\widetilde{C}|) - 3$, *where* $\mathfrak{C}$ *is the family of all virtual minimal cycles of* G.

Proof. Let $C \in \mathfrak{C}^v$, thanks to definition of $\mathfrak{C}^{\mathrm{ls}}_{\Pi^*}(C)$, $\widetilde{C} \in \mathfrak{C}^{\mathrm{ls}}_{\Pi}(C)$. By proposition 4.16 $|\widetilde{C}| = \min_{C'\in\mathfrak{C}^{\mathrm{ls}}_{\Pi^*}(C)}|C'|$. Thus, from the theorems 5.9 and 4.17

$$p_2(\mathcal{B}_{\overline{\Gamma}}) \le \min_{C\in\mathfrak{C}^v}(|\widetilde{C}|) - 3 = \min_{C\in\mathfrak{C}^v}(\min_{C'\in\mathfrak{C}^{\mathrm{ls}}_{\Pi^*}(C)}|C'|) - 3 \ge p_2(\mathcal{B}_{\overline{\Gamma}}).$$

So $p_2(\mathcal{B}_{\overline{\Gamma}}) = \min_{C\in\mathfrak{C}^v}(|\widetilde{C}|) - 3$.

■

5.11 Example. Let G be the graph of the figure 5 and $\Gamma = \Gamma(G)$ the clique complex generated by G. The subgraph C of G whose edge-set $E(C) = \{\{a,c\},\{c,d\},\{d,e\},\{e,a\}\}$ is the only minimal cycle of G. We associate to the facets $F_1 = \{a,b,c\}$ and $F_2 = \{c,d\}$ the following matrices:

$$M_{F_1} = \begin{pmatrix} a & z \\ z & c \end{pmatrix} \text{ and } M_{F_2} = \begin{pmatrix} e & w & x \\ w & x & d \end{pmatrix}.$$

Thus the graph $\overline{G}'$ of the figure 5 is the 1-skeleton of $\overline{\Gamma}^{\Pi^*} = \overline{\Gamma}'$.



Figure 5:

Besides, by a proposition of the preprint [4], $\mathcal{J}_{\overline{\Gamma}}$ is a toric ideal. Moreover, we see that the conditions of the theorem 5.10 required to Γ and its extension $\overline{\Gamma}$ are satisfied, so:

$$\begin{aligned} p_2(\mathcal{B}_{\overline{\Gamma}}) &= p_2(I_{\overline{\Gamma}^{\Pi^*}}) \\ &= (|C| + |R_1^{(\Pi^*)}(C)| + \textstyle\sum_{e\in R_3^{(\Pi^*)}(C)} \eta_e) - 3 \\ &= (4+1+2) - 3 = 4. \end{aligned}$$

References

[1] W. Adams, P. Loustaunau, *An introduction to Gröbner Bases*, Graduate studies in mathematics, American Mathematical Society.

[2] W. Bruns, J. Herzog, *Semigroup ring and simplicial complexes*, J. of Pure and App. Math, **122** (1997), 185-208.

[3] H. de Alba Casillas, Nombres de Betti d'idéaux binomiaux, P.H.D. thesis.

[4] H. de Alba Casillas, M. Morales, *Sum of toric ideals*, preprint.

[5] D. Eisenbud, *Commutative algebra with a view toward algebraic geometry*, Graduate Text in Mathematics, Springer, 1995.

[6] D. Eisenbud, S. Goto, *Linear free resolutions and minimal multiplicity*, J. Algebra **88** (1984) 89-133.

[7] D. Eisenbud, B. Sturmfels, *Binomial Ideals*, Duke Math. J., **84** (1996), 1-45.

[8] D. Eisenbud, M. Green, K. Hulek, S. Popescu, *Restricting linear syzygies algebra and geometry*, Composition Math. **141** (2005), 1460-1478.

[9] O. Fernández-Ramos, P. Gimenez, *First nonlinear syzygies of ideals associated to graphs*, Comm Algebra **37** (2009), no. 6, 1921-1933.

[10] R. Fröberg, *On Stanley-Reisner rings*, Topics in Algebra, Banach Center Publications, vol.26, part 2 (1990), 57-70.

[11] J. Harris, *Algebraic geometry a first course*, Springer, 1992.

[12] J. Herzog, M. Kühl, *On the Betti numbers of finite pure linear resolutions*, Comm. algebra **12** (1984), no. 13-14,1627-1646.

[13] J. Herzog, T. Hibi, *Monomial ideals*, Graduate Texts in Mathematics, Springer, 2011.

[14] M., Hochster, *Cohen-Macaulay rings, combinatorics and simplicial complexes*, in "Ring Theory II", pp. 171-223, Dekker, New York, 1977

[15] M. L. Ha, M. Morales, *Binomial extension of simplicial ideals*, http://arxiv.org/abs/0911.5423.

[16] J. Munkres, *Elements of algebraic Topoloy*, Addison-Wesley, 1984.

[17] M. Morales, *Simplicial ideals, 2-linear ideals and arithmetical rank*, J. Algebra (2010), doi:10.1016/j.algebra.2010.08.025.

[18] H. Lam, M. Morales, *Fiber cones of codimension 2 lattice ideals*, Com. Alg., **37** (2009), 1-31.

[19] E. Mïller, B. Sturmfels, *Combinatorial commutative algebra*, Springer, 2005.

[20] E. Spanier, Algebraic Topology, McGrawHill, Inc., 1966.

[21] B. Sturmfels, *Gröbner bases and convex polytopes*, America Mathematical Society, 1996.